

# EQUIVALENCE OF HADAMARD MATRICES

BY  
W. D. WALLIS AND JENNIFER WALLIS

## ABSTRACT

Suppose  $m$  is a square-free odd integer, and  $A$  and  $B$  are any two Hadamard matrices of order  $4m$ . We will show that  $A$  and  $B$  are equivalent over the integers (that is,  $B$  can be obtained from  $A$  using elementary row and column operations which involve only integers).

**Integral equivalence.** If  $A$  and  $B$  are matrices over the ring  $\mathbf{Z}$  of integers,  $A$  and  $B$  are called *equivalent* ( $A \sim B$ ) if there are  $\mathbf{Z}$ -matrices  $P$  and  $Q$ , of determinant  $\pm 1$ , such that

$$B = P \cdot A \cdot Q.$$

This is the same as saying that  $B$  can be obtained from  $A$  by performing some sequence of the following operations:

- (a) add an integer multiple of one row to another,
- (b) negate some row,
- (c) reorder the rows,

and the corresponding column operations. The main result about equivalence is

**LEMMA.** *If  $A$  is any  $n \times n$   $\mathbf{Z}$ -matrix, then there is a unique  $\mathbf{Z}$ -matrix*

$$D = \text{diag}(a_1, a_2, \dots, a_n)$$

*such that  $A \sim D$  and*

$$a_1 \mid a_2 \mid \dots \mid a_r, \quad a_{r+1} = \dots = a_n = 0,$$

---

Received January 29, 1969.

where the  $a_i$  are non-negative. The greatest common divisor of  $i \times i$  subdeterminants of  $A$  is

$$a_1 a_2 a_3 \cdots a_i.$$

If  $A \sim E$  where

$$E = \left[ \begin{array}{ccc|c} a_1 & & & 0 \\ & a_2 & & \\ & & \cdots & \\ & & & a_i \\ \hline & & & 0 \\ & & & F \end{array} \right]$$

then  $a_{i+1}$  is the greatest common divisor of non-zero elements of  $F$ .

The  $a_i$  are called *invariants* of  $A$ .

**Hadamard matrices.** An Hadamard matrix  $A$  of order  $n$  is an  $n \times n$  matrix whose elements are  $\pm 1$  and which satisfies

$$AA^T = nI_n.$$

(See, for example, Chapter 14 of [1]). If  $A$  is any Hadamard matrix we can find an Hadamard matrix  $H$  satisfying

$$H \sim A,$$

$$H = \left[ \begin{array}{c|cccc} 1 & 1 & 1 & \cdots & 1 \\ \hline 1 & & & & \\ 1 & & & & \\ \vdots & & & & \\ 1 & & & & \end{array} \right] \begin{array}{c} \\ \\ \\ B \\ \end{array}$$

simply by negating rows and columns,  $H$  is then *normalized*.

The determinant of an Hadamard matrix is

$$\pm n^{1/2n}$$

**Certain invariants.** Suppose  $A$  is an Hadamard matrix of order  $n = 4m$ . We will find some of the invariants of  $A$ . There is no loss of generality in assuming that  $A$  is normalized.

Since every element is  $\pm 1$ ,  $a_1$  must be 1. Now subtract the first row from every other row, and then the first column from every other column. The resulting matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & K \end{bmatrix}$$

is equivalent to  $A$ , and every element of  $K$  is 0 or  $-2$ . So

$$a_2 = 2.$$

By definition

$$a_{4m} = \pm \frac{|A|}{a_1 a_2 \cdots a_{4m-1}};$$

the numerator is  $(4m)^{2m}$ , and the denominator is the greatest common divisor of the  $(4m-1)$ -subdeterminants of  $A$ . We shall now evaluate this greatest common divisor.

Suppose  $C$  is any  $(4m-1)$ -subdeterminant of  $A$ . Then

$$\begin{aligned} A &\sim \left[ \begin{array}{c|c} \pm 1 & \pm 1 \cdots \pm 1 \\ \hline \pm 1 & \\ \dots & \\ \pm 1 & \end{array} \right] \\ &\sim \left[ \begin{array}{c|c} 1 & 1 \cdots 1 \\ \hline 1 & \\ \dots & \\ 1 & \end{array} \right] = F; \end{aligned}$$

$B$  is obtained from  $C$  by negating rows and columns, hence

$$|B| = \pm |C|.$$

$F$  is Hadamard, so

$$FF^T = 4mI_{4m};$$

but

$$FF^T = \left[ \begin{array}{c|c} 4m & \\ \hline & BB^T + J_{4m-1} \end{array} \right]$$

where  $J_v$  is the  $v \times v$  matrix whose every element is  $+1$ . Therefore

$$BB^T = 4mI_{4m-1} - J_{4m-1}.$$

$$|(r-\lambda)I_v + \lambda J_v| = \{r + (v-1)\lambda\} (r-\lambda)^{v-1}$$

[2, p. 99], whence, putting  $v = r = 4m - 1$ ,  $\lambda = -1$ ,

$$\begin{aligned} |B|^2 &= (4m)^{4m-2}, \\ |C| &= \pm (4m)^{2m-1}. \end{aligned}$$

This works for any  $(4m-1)$ -subdeterminant, so the greatest common divisor is  $(4m)^{2m-1}$ , and

$$a_{4m} = 4m.$$

**When  $m$  is odd and square-free.** We continue the notation of the last section, and further suppose that  $m$  is odd and square-free. Since 2 must divide every invariant but  $a_1$ , write

$$b_i = \frac{1}{2}a_i, \quad i > 1.$$

$$|A| = \pm (4m)^{2m} = \pm 2^{4m}m^{2m};$$

but on the other hand

$$\begin{aligned} |A| &= \pm \prod a_i \\ &= \pm 2^{4m}m \prod_{i=2}^{4m-1} b_i; \end{aligned}$$

therefore

$$\prod_{i=2}^{4m-1} b_i = m^{2m-1}.$$

If  $p$  is any prime factor of  $m$ , then  $p^{2m-1}$  is a factor of this product.  $p^2$  does not divide  $a_{4m}$ , so  $p^2$  cannot divide any of the  $b_i$ . Hence exactly  $2m - 1$  of them must have a factor  $p$ . By the property

$$a_1 | a_2 | a_3 \cdots,$$

these must be  $b_{2m+1}, \dots, b_{4m-1}$ . Hence  $m$  divides each of these  $b_i$ ; the rest must all be 1. We have

**THEOREM 1.** *If  $A$  is Hadamard of order  $4m$ , where  $m$  is odd and square-free then the invariants of  $A$  are*

1 (once)  
 2 ( $2m - 1$  times)  
 $2m$  ( $2m - 1$  times)  
 $4m$  (once).

**COROLLARY.** *Any two Hadamard matrices of order  $4m$ , where  $m$  is and odd square-free, are  $\mathbf{Z}$ -equivalent.*

**When  $m$  is even.** We can partially extend Theorem 1 to the case where  $m$  is even and square-free. If  $H$  is an Hadamard matrix of order  $2m$ , then

$$A = \begin{bmatrix} H & H \\ H & -H \end{bmatrix}$$

is Hadamard of order  $4m$ . Now

$$\begin{aligned} A &\sim \begin{bmatrix} H & 0 \\ 0 & -2H \end{bmatrix} \\ &\sim \begin{bmatrix} D & 0 \\ 0 & 2D \end{bmatrix}, \end{aligned}$$

where  $D$  is the diagonal matrix of Theorem 1 corresponding to  $H$ . (The theorem can be applied, as  $\frac{1}{2}m$  is odd). Thus  $A$  is equivalent to a diagonal matrix with elements

1 (once)  
 2 ( $m$  times)  
 $m$  ( $m - 1$  times)  
 $2m$  ( $m$  times)  
 4 ( $m - 1$  times)  
 $4m$  (once).

There is a  $(2m)$ -subdeterminant

$$1 \cdot 2^m \cdot m^{m-1} = 2^{2m-1}k,$$

where  $k$  is odd, and another

$$1 \cdot 2^m \cdot 4^{m-1} = 2^{3m-2}.$$

The greatest common divisor of these is  $2^{2m-1}$ , so

$$a_1 a_2 \cdots a_{2m} \leq 2^{2m-1}.$$

On the other hand each  $a_i$  (after  $a_1$ ) is divisible by 2, hence

$$a_1 a_2 \cdots a_{2m} \geq 2^{2m-1};$$

equality holds, and

$$a_1 = 1, a_2 = a_3 = \cdots = a_{2m} = 2.$$

Now we find  $a_{4m-1}$ . From an earlier result

$$a_1 a_2 \cdots a_{4m-1} = (4m)^{2m-1}.$$

One  $(4m-2)$ -subdeterminant is

$$\delta = 2(4m)^{2m-2}$$

obtained by deleting the diagonal elements  $4m$  and  $2m$ . Every other  $(4m-2)$ -subdeterminant results from replacing one or two of the diagonal elements of  $\delta$

by  $2m$  or  $4m$  (or both); every diagonal element of  $\delta$  divides  $2m$ , so  $\delta$  divides every other  $(4m-2)$ -subdeterminant. Therefore

$$a_1 a_2 \cdots a_{4m-2} = 2(4m)^{2m-2},$$

$$a_{4m-1} = 2m.$$

Since  $m$  is square-free,

$$a_{2m+1} = a_{2m+2} = \cdots = a_{4m-2} = 2m.$$

Thus we have proven

**THEOREM 2.** *If  $m$  is even and square-free, and if there is an Hadamard matrix of order  $2m$ , then there is an Hadamard matrix of order  $4m$  of the type in Theorem 1.*

However it is possible that there are also matrices of these orders with other invariants.

**Trivial cases.** In the trivial cases ( $n = 1$  or  $2$ ) the invariants are of the type in Theorem 1.

**A matrix of order 16.** Finally we show that there is an Hadamard matrix whose invariants are not in the form of Theorem 1. Let  $H$  be an Hadamard matrix of order 4. The invariants of  $H$  are thus  $\{1, 2, 2, 4\}$ . If  $A$  is the direct product  $H \times H$  then

$$A \sim \text{diag}(1, 2, 2, 4) \times \text{diag}(1, 2, 2, 4).$$

This is a diagonal matrix with elements

1 (once)  
2 (four times)  
4 (six times)  
8 (four times)  
16 (once),

and these are clearly the invariants of  $A$ .

#### REFERENCES

1. M. Hall Jr., *Combinatorial Theory*, Blaisdell, Waltham, Mass., 1967.
2. H. J. Ryser, *Combinatorial Mathematics*, (Carus Monograph No. 14), Wiley, New York, 1963.

LA TROBE UNIVERSITY,  
BUNDOORA,  
VICTORIA, 3083,  
AUSTRALIA