

On Full Orthogonal Designs in Order 72

S. Georgiou*, C. Koukouvinos*, and Jennifer Seberry†

July 23, 2000

Abstract

We find new full orthogonal designs in order 72 and show that of 2700 possible $OD(72; s_1, s_2, s_3, 72 - s_1 - s_2 - s_3)$ 335 are known, of 432 possible $OD(72; s_1, s_2, 72 - s_1 - s_2)$ 308 are known. All possible $OD(72; s_1, 72 - s_1)$ are known.

Key words and phrases: Construction, sequences, circulant matrices, amicable sets, orthogonal designs.

AMS Subject Classification: Primary 05B15, 05B20, Secondary 62K05.

1 Introduction

An *orthogonal design* of order n and type (s_1, s_2, \dots, s_u) ($s_i > 0$), denoted $OD(n; s_1, s_2, \dots, s_u)$, on the commuting variables x_1, x_2, \dots, x_u is an $n \times n$ matrix A with entries from $\{0, \pm x_1, \pm x_2, \dots, \pm x_u\}$ such that

$$AA^T = \left(\sum_{i=1}^u s_i x_i^2 \right) I_n$$

Alternatively, the rows of A are formally orthogonal and each row has precisely s_i entries of the type $\pm x_i$. In [1], where this was first defined, it was mentioned that

$$A^T A = \left(\sum_{i=1}^u s_i x_i^2 \right) I_n$$

and so our alternative description of A applies equally well to the columns of A . It was also shown in [1] that $u \leq \rho(n)$, where $\rho(n)$ (Radon's function) is defined by $\rho(n) = 8c + 2^d$, when $n = 2^a b$, b odd, $a = 4c + d$, $0 \leq d < 4$.

A weighing matrix $W = W(n, k)$ is a square matrix with entries $0, \pm 1$ having k non-zero entries per row and column and inner product of distinct rows zero. Hence W satisfies $WW^T = kI_n$, and W is equivalent to an orthogonal design $OD(n; k)$. The number k is called the *weight* of W . If $k = n$, that is, all the entries of W are ± 1 and $WW^T = nI_n$, then W is called an Hadamard matrix of order n . In this case $n = 1, 2$ or $n \equiv 0 \pmod{4}$.

*Department of Mathematics, National Technical University of Athens, Zografou 15773, Athens, Greece.

†Department of Computer Science, University of Wollongong, Wollongong, NSW, 2522, Australia.

Given the sequence $A = \{a_1, a_2, \dots, a_n\}$ of length n the *non-periodic autocorrelation function* $N_A(s)$ is defined as

$$N_A(s) = \sum_{i=1}^{n-s} a_i a_{i+s}, \quad s = 0, 1, \dots, n-1, \quad (1)$$

If $A(z) = a_1 + a_2 z + \dots + a_n z^{n-1}$ is the associated polynomial of the sequence A , then

$$A(z)A(z^{-1}) = \sum_{i=1}^n \sum_{j=1}^n a_i a_j z^{i-j} = N_A(0) + \sum_{s=1}^{n-1} N_A(s)(z^s + z^{-s}), \quad z \neq 0. \quad (2)$$

Given A as above of length n the *periodic autocorrelation function* $P_A(s)$ is defined, reducing $i + s$ modulo n , as

$$P_A(s) = \sum_{i=1}^n a_i a_{i+s}, \quad s = 0, 1, \dots, n-1. \quad (3)$$

The following theorem which uses four circulant matrices in the Goethals-Seidel array is very useful in our construction for orthogonal designs.

Theorem 1 [2, Theorem 4.49] *Suppose there exist four circulant matrices A, B, C, D of order n satisfying*

$$AA^T + BB^T + CC^T + DD^T = fI_n$$

Let R be the back diagonal matrix. Then

$$GS = \begin{pmatrix} A & BR & CR & DR \\ -BR & A & D^T R & -C^T R \\ -CR & -D^T R & A & B^T R \\ -DR & C^T R & -B^T R & A \end{pmatrix}$$

is a $W(4n, f)$ when A, B, C, D are $(0, 1, -1)$ matrices, and an orthogonal design $OD(4n; s_1, s_2, \dots, s_u)$ on x_1, x_2, \dots, x_u when A, B, C, D have entries from $\{0, \pm x_1, \dots, \pm x_u\}$ and $f = \sum_{j=1}^u (s_j x_j^2)$. \square

Corollary 1 *If there are four sequences A, B, C, D of length n with entries from $\{0, \pm x_1, \pm x_2, \pm x_3, \pm x_4\}$ with zero periodic or non-periodic autocorrelation function, then these sequences can be used as the first rows of circulant matrices which can be used in the Goethals-Seidel array to form an $OD(4n; s_1, s_2, s_3, s_4)$. We note that if the non-periodic autocorrelation function is zero, then there are sequences of length $n + m$ for all $m \geq 0$. \square*

This method for constructing orthogonal designs was used in [3, 5, 6]. Furthermore, in [6] a complete study for two variable orthogonal designs in order 36 was given, and in [3] a complete study for three and four variable orthogonal designs in order 36 was also presented.

Throughout this paper we will use the definition and notation of Koukouvinos, Mitrouli, Seberry and Karabelas [5].

A pair of matrices A, B is said to be amicable (anti-amicable) if $AB^T - BA^T = 0$ ($AB^T + BA^T = 0$). Following [4] a set $\{A_1, A_2, \dots, A_{2n}\}$ of square real matrices is said to be *amicable* if

$$\sum_{i=1}^n (A_{\sigma(2i-1)} A_{\sigma(2i)}^T - A_{\sigma(2i)} A_{\sigma(2i-1)}^T) = 0 \quad (4)$$

for some permutation σ of the set $\{1, 2, \dots, 2n\}$. For simplicity, we will always take $\sigma(i) = i$ unless otherwise specified. So

$$\sum_{i=1}^n (A_{2i-1}A_{2i}^T - A_{2i}A_{2i-1}^T) = 0. \quad (5)$$

Clearly a set of mutually amicable matrices is amicable, but the converse is not true in general. Throughout this paper R_k denotes the back diagonal identity matrix of order k .

Let $\{A_i\}_{i=1}^8$ be an amicable set of circulant matrices of order t , satisfying the additive property for (s_1, s_2, \dots, s_k) . Then the Kharaghani array

$$H = \begin{pmatrix} A_1 & A_2 & A_4R_n & A_3R_n & A_6R_n & A_5R_n & A_8R_n & A_7R_n \\ -A_2 & A_1 & A_3R_n & -A_4R_n & A_5R_n & -A_6R_n & A_7R_n & -A_8R_n \\ -A_4R_n & -A_3R_n & A_1 & A_2 & -A_8^T R_n & A_7^T R_n & A_6^T R_n & -A_5^T R_n \\ -A_3R_n & A_4R_n & -A_2 & A_1 & A_7^T R_n & A_8^T R_n & -A_5^T R_n & -A_6^T R_n \\ -A_6R_n & -A_5R_n & A_8^T R_n & -A_7^T R_n & A_1 & A_2 & -A_4^T R_n & A_3^T R_n \\ -A_5R_n & A_6R_n & -A_7^T R_n & -A_8^T R_n & -A_2 & A_1 & A_3^T R_n & A_4^T R_n \\ -A_8R_n & -A_7R_n & -A_6^T R_n & A_5^T R_n & A_4^T R_n & -A_3^T R_n & A_1 & A_2 \\ -A_7R_n & A_8R_n & A_5^T R_n & A_6^T R_n & -A_3^T R_n & -A_4^T R_n & -A_2 & A_1 \end{pmatrix}$$

is an $OD(8t; s_1, s_2, \dots, s_k)$.

The Kharaghani array which uses amicable sets of eight matrices is also very useful in our constructions for orthogonal designs.

The following lemma applies a lemma given in Georgiou, Koukouvinos, Mitrouli and Seberry [3] to determine the number of possible tuples to be searched determining the size of search space for orthogonal designs in order 72.

Lemma 1 *Let $n = 4m = 72$ be the order of an orthogonal design then the number of cases which must be studied to determine whether all orthogonal designs exist is*

- (i) $\frac{1}{4}n^2 = 1296$ when 2-tuples are considered;
- (ii) $\frac{n}{72}(2n^2 + 3n - 6) = 10578$ when 3-tuples are considered;
- (iii) $\frac{1}{576}(n^4 + 6n^3 - 2n^2 - 24n) = 50523$ when 4-tuples are considered.

2 New full orthogonal designs from smaller orders

Theorem 2 *There are $OD(72; s_1, s_1, 36-s_1, 36-s_1)$ constructed using the full $OD(36; s_1, 36-s_1)$ given in [3, Theorem8] for:*

$$\begin{array}{cccc} (1, 1, 35, 35) & (6, 6, 30, 30) & (11, 11, 25, 25) & (15, 15, 21, 21) \\ (2, 2, 34, 34) & (7, 7, 29, 29) & (12, 12, 24, 24) & (16, 16, 20, 20) \\ (3, 3, 33, 33) & (8, 8, 28, 28) & (13, 13, 23, 23) & (17, 17, 19, 19) \\ (4, 4, 32, 32) & (9, 9, 27, 27) & (14, 14, 22, 22) & (18, 18, 18, 18) \\ (5, 5, 31, 31) & (10, 10, 26, 26) & & \end{array}$$

Proof. We use the amicable orthogonal designs of type $AOD(2; (1, 1), (1, 1))$ in order two with the two variable designs in order 36 to obtain the desired designs in order 72. \square

Theorem 3 *There are full $OD(72; s_1, s_2, s_3, 72 - s_1 - s_2 - s_3)$ constructed using the full $OD(36; s_1, s_2, 36 - s_1 - s_2)$ and $OD(36; s_1, s_2, s_3, 36 - s_1 - s_2 - s_3)$ designs in order 36 for the 4-tuples given in Table 1.*

Theorem 4 *There are $OD(72; s_1, s_1, 2s_2, 2s_3, 72 - 2s_1 - 2s_2 - 2s_3)$ constructed using the Multiplication Theorem [2, Lemma 4.11] with the full $OD(36; s_1, s_2, s_3, 36 - s_1 - s_2 - s_3)$ given in [3, Theorem2] for the values given in Table 2.*

Theorem 5 *There are $OD(72; s_1, \dots, s_u, 72 - \sum_{i=1}^u s_i)$ constructed using $OD(24; t_1, t_2, t_3, t_4, t_5, t_6, t_7, 24 - \sum_{i=1}^7 t_i)$ as follows;*

1, 2, 9, 10, 50	2, 10, 15, 45	3, 6, 6, 27, 30	4, 8, 20, 40
1, 5, 6, 10, 50	2, 9, 27, 34	3, 6, 15, 18, 30	4, 20, 24, 24
2, 2, 8, 10, 50	2, 17, 53	3, 6, 18, 18, 27	5, 6, 6, 55
2, 7, 63	3, 6, 6, 9, 48	3, 15, 18, 18, 18	8, 12, 12, 40

Proof. We use circulant and back circulant matrices to replace the variables in orthogonal designs of order 24 in the cases of table 3. If the matrix is symmetric such as that with first row $\mathbf{c} \ \mathbf{b} \ \mathbf{b}$; we use the circulant matrix. However if the matrix would not be symmetric such as that with first row $\mathbf{a} \ \mathbf{b} \ \mathbf{b}$ we use the back circulant matrix to replace the variables in the orthogonal design.

Table 5 gives the results for $(4, 8, 20, 40)$, $(4, 20, 24, 24)$ and $(8, 12, 12, 40)$ using the symmetric matrix $C = W(6, 5)$ and the identity matrix I with the variables x, y, z and w to replace the variables of an orthogonal design of order 12. \square

In table 6 we present the new amicable sets of eight matrices which can be used in the Kharaghani array to construct some new full orthogonal designs in order 72.

We use the results of the theorem 5 to form Table 8.

3 Full designs with even parameters

We note that Seberry [7] showed that if all $OD(n; x, y, n-x-y)$ exist then all $OD(2n; z, w, 2n-z-w)$ exist for $s \geq 0$ an integer. In particular if all $OD(2^t p; x, y, 2^t p - x - y)$ exist, for some odd integer p , then all $OD(2^{t+s} p; z, w, 2^{t+s} p - z - w)$ exist for $s \geq 0$ an integer we observe

Lemma 2 *If all $OD(2^t p; 2x, 2y, 2^t p - 2x - 2y)$ exist, for some odd integer p , then all $OD(2^{t+s} p; 2z, 2w, 2^{t+s} p - 2z - 2w)$ exist for $s \geq 0$ an integer.*

Corollary 2 *If $OD(72; 12, 22, 38)$, $OD(72; 14, 20, 38)$ and $OD(72; 14, 28, 30)$ exist then all $OD(2^{s+3} 9; 2z, 2w, 2^{s+3} 9 - 2z - 2w)$ exist for $s \geq 0$ an integer.*

Proof. A search of full $OD(72; x, y, 72 - x - y)$ show only the parameters indicated are as yet unsolved. \square

(1, 1, 2, 68)	(2, 4, 12, 54)	(3, 3, 16, 50)	(4, 21, 21, 26)	(8, 10, 18, 36)
(1, 1, 4, 66)	(2, 4, 22, 44)	(3, 3, 18, 48)	(4, 22, 22, 24)	(8, 12, 16, 36)
(1, 1, 6, 64)	(2, 4, 27, 39)	(3, 3, 20, 46)	(4, 22, 23, 23)	(8, 12, 26, 26)
(1, 1, 10, 60)	(2, 4, 32, 34)	(3, 3, 22, 44)	(5, 5, 10, 52)	(8, 16, 16, 32)
(1, 1, 12, 58)	(2, 4, 33, 33)	(3, 3, 30, 36)	(5, 5, 12, 50)	(8, 16, 20, 28)
(1, 1, 16, 54)	(2, 5, 5, 60)	(3, 4, 12, 53)	(5, 5, 16, 46)	(8, 16, 24, 24)
(1, 1, 18, 52)	(2, 5, 10, 55)	(3, 4, 15, 50)	(5, 5, 26, 36)	(8, 20, 20, 24)
(1, 1, 20, 50)	(2, 5, 15, 50)	(3, 5, 16, 48)	(5, 7, 10, 50)	(9, 9, 12, 42)
(1, 1, 22, 48)	(2, 5, 6, 58)	(3, 6, 6, 57)	(5, 10, 16, 41)	(9, 9, 16, 38)
(1, 1, 26, 44)	(2, 6, 8, 56)	(3, 6, 9, 54)	(5, 10, 21, 36)	(9, 9, 18, 36)
(1, 1, 28, 42)	(2, 6, 10, 54)	(3, 6, 12, 51)	(5, 10, 26, 31)	(9, 9, 26, 28)
(1, 1, 34, 36)	(2, 6, 12, 52)	(3, 6, 15, 48)	(5, 15, 16, 36)	(9, 11, 16, 36)
(1, 2, 4, 65)	(2, 6, 14, 50)	(3, 6, 30, 33)	(5, 15, 26, 26)	(9, 12, 15, 36)
(1, 2, 5, 64)	(2, 6, 16, 48)	(3, 7, 12, 50)	(6, 6, 6, 54)	(9, 18, 18, 27)
(1, 2, 18, 51)	(2, 6, 24, 40)	(3, 9, 12, 48)	(6, 6, 10, 50)	(10, 10, 13, 39)
(1, 2, 19, 50)	(2, 6, 32, 32)	(3, 9, 30, 30)	(6, 6, 12, 48)	(10, 10, 16, 36)
(1, 2, 34, 35)	(2, 8, 8, 54)	(3, 12, 18, 39)	(6, 6, 14, 46)	(10, 10, 18, 34)
(1, 3, 4, 64)	(2, 8, 12, 50)	(3, 12, 21, 36)	(6, 6, 15, 45)	(10, 10, 20, 32)
(1, 3, 18, 50)	(2, 8, 14, 48)	(3, 15, 18, 36)	(6, 6, 18, 42)	(10, 10, 22, 30)
(1, 3, 34, 34)	(2, 8, 18, 44)	(4, 4, 16, 48)	(6, 6, 24, 36)	(10, 10, 25, 27)
(1, 4, 12, 55)	(2, 8, 26, 36)	(4, 4, 32, 32)	(6, 6, 27, 33)	(10, 10, 26, 26)
(1, 4, 13, 54)	(2, 9, 9, 52)	(4, 6, 6, 56)	(6, 6, 28, 32)	(10, 12, 25, 25)
(1, 4, 22, 45)	(2, 9, 11, 50)	(4, 6, 8, 54)	(6, 6, 30, 30)	(10, 13, 13, 36)
(1, 4, 23, 44)	(2, 9, 16, 45)	(4, 6, 12, 50)	(6, 8, 8, 50)	(10, 13, 23, 26)
(1, 5, 12, 54)	(2, 9, 25, 36)	(4, 6, 24, 38)	(6, 8, 10, 48)	(10, 16, 16, 30)
(1, 5, 22, 44)	(2, 10, 10, 50)	(4, 6, 25, 37)	(6, 9, 9, 48)	(10, 16, 18, 28)
(1, 6, 16, 49)	(2, 10, 25, 35)	(4, 6, 30, 32)	(6, 9, 12, 45)	(10, 16, 20, 26)
(1, 6, 17, 48)	(2, 10, 30, 30)	(4, 6, 31, 31)	(6, 9, 21, 36)	(10, 16, 23, 23)
(1, 7, 16, 48)	(2, 11, 11, 48)	(4, 7, 7, 54)	(6, 10, 10, 46)	(10, 18, 18, 26)
(1, 10, 10, 51)	(2, 11, 15, 44)	(4, 8, 8, 52)	(6, 10, 24, 32)	(11, 11, 20, 30)
(1, 10, 11, 50)	(2, 12, 24, 34)	(4, 8, 16, 44)	(6, 11, 11, 44)	(12, 12, 12, 36)
(1, 16, 18, 37)	(2, 12, 26, 32)	(4, 8, 24, 36)	(6, 12, 18, 36)	(12, 12, 16, 32)
(1, 16, 19, 36)	(2, 12, 27, 31)	(4, 8, 26, 34)	(6, 12, 24, 30)	(12, 12, 24, 24)
(1, 17, 18, 36)	(2, 12, 29, 29)	(4, 9, 9, 50)	(6, 12, 25, 29)	(12, 14, 14, 32)
(2, 2, 2, 66)	(2, 13, 13, 44)	(4, 11, 11, 46)	(6, 12, 27, 27)	(12, 14, 23, 23)
(2, 2, 4, 64)	(2, 14, 14, 42)	(4, 11, 13, 44)	(6, 15, 15, 36)	(12, 15, 15, 30)
(2, 2, 6, 62)	(2, 14, 24, 32)	(4, 12, 12, 44)	(6, 15, 21, 30)	(12, 16, 16, 28)
(2, 2, 9, 59)	(2, 16, 16, 38)	(4, 12, 16, 40)	(6, 16, 24, 26)	(12, 16, 20, 24)
(2, 2, 12, 56)	(2, 16, 18, 36)	(4, 12, 24, 32)	(6, 16, 25, 25)	(12, 18, 18, 24)
(2, 2, 14, 54)	(2, 16, 24, 30)	(4, 12, 25, 31)	(6, 18, 18, 30)	(12, 18, 21, 21)
(2, 2, 16, 52)	(2, 16, 26, 28)	(4, 12, 27, 29)	(6, 18, 24, 24)	(13, 13, 16, 30)
(2, 2, 17, 51)	(2, 16, 27, 27)	(4, 12, 28, 28)	(6, 20, 23, 23)	(13, 13, 18, 28)
(2, 2, 18, 50)	(2, 17, 17, 36)	(4, 13, 13, 42)	(6, 22, 22, 22)	(13, 13, 20, 26)
(2, 2, 22, 46)	(2, 17, 19, 34)	(4, 13, 16, 39)	(7, 7, 12, 46)	(13, 16, 17, 26)
(2, 2, 24, 44)	(2, 18, 18, 34)	(4, 13, 26, 29)	(7, 7, 16, 42)	(14, 14, 18, 26)
(2, 2, 25, 43)	(2, 18, 25, 27)	(4, 14, 27, 27)	(8, 8, 8, 48)	(14, 16, 21, 21)
(2, 2, 26, 42)	(2, 18, 26, 26)	(4, 16, 16, 36)	(8, 8, 10, 46)	(15, 15, 16, 26)
(2, 2, 32, 36)	(2, 20, 25, 25)	(4, 16, 18, 34)	(8, 8, 14, 42)	(15, 15, 20, 22)
(2, 2, 34, 34)	(2, 21, 21, 28)	(4, 16, 20, 32)	(8, 8, 16, 40)	(16, 16, 16, 24)
(2, 3, 3, 64)	(2, 22, 22, 26)	(4, 16, 24, 28)	(8, 8, 18, 38)	(16, 16, 18, 22)
(2, 3, 16, 51)	(2, 22, 24, 24)	(4, 16, 26, 26)	(8, 8, 20, 36)	(16, 16, 20, 20)
(2, 3, 19, 48)	(3, 3, 4, 62)	(4, 17, 17, 34)	(8, 8, 24, 32)	(16, 18, 18, 20)
(2, 4, 6, 60)	(3, 3, 6, 60)	(4, 18, 18, 32)	(8, 8, 26, 30)	(16, 18, 19, 19)
(2, 4, 11, 55)	(3, 3, 12, 54)	(4, 18, 25, 25)	(8, 10, 10, 44)	(18, 18, 18, 18)

Table 1: Full 4-variable $OD(72; s_1, s_2, s_3, 72 - s_1 - s_2 - s_3)$ constructed from full three and four variable designs in order 36

(1, 1, 2, 4, 64)	(2, 2, 16, 26, 26)	(3, 3, 6, 6, 54)	(6, 6, 6, 6, 48)
(1, 1, 2, 18, 50)	(2, 2, 17, 17, 34)	(3, 3, 6, 12, 48)	(6, 6, 6, 18, 36)
(1, 1, 2, 34, 34)	(2, 2, 18, 25, 25)	(3, 3, 6, 30, 30)	(6, 6, 6, 27, 27)
(1, 1, 4, 12, 54)	(2, 3, 3, 16, 48)	(3, 3, 12, 18, 36)	(6, 6, 12, 24, 24)
(1, 1, 4, 22, 44)	(2, 4, 6, 6, 54)	(4, 4, 16, 16, 32)	(6, 6, 15, 15, 30)
(1, 1, 6, 16, 48)	(2, 4, 11, 11, 44)	(4, 6, 6, 6, 50)	(6, 9, 9, 12, 36)
(1, 1, 10, 10, 50)	(2, 4, 12, 27, 27)	(4, 6, 6, 24, 32)	(6, 12, 18, 18, 18)
(1, 1, 16, 18, 36)	(2, 4, 22, 22, 22)	(4, 6, 12, 25, 25)	(8, 8, 8, 16, 32)
(2, 2, 2, 2, 64)	(2, 5, 5, 10, 50)	(4, 8, 8, 16, 36)	(8, 8, 10, 10, 36)
(2, 2, 2, 12, 54)	(2, 6, 8, 8, 48)	(4, 8, 8, 26, 26)	(8, 8, 16, 20, 20)
(2, 2, 2, 22, 44)	(2, 6, 16, 24, 24)	(4, 12, 12, 12, 32)	(8, 16, 16, 16, 16)
(2, 2, 4, 32, 32)	(2, 8, 8, 18, 36)	(4, 12, 16, 16, 24)	(9, 9, 18, 18, 18)
(2, 2, 6, 12, 50)	(2, 9, 9, 16, 36)	(4, 13, 13, 16, 26)	(10, 10, 13, 13, 26)
(2, 2, 9, 9, 50)	(2, 10, 10, 25, 25)	(4, 16, 16, 18, 18)	(10, 10, 16, 16, 20)
(2, 2, 12, 24, 32)	(2, 16, 18, 18, 18)	(5, 5, 10, 16, 36)	(10, 10, 16, 18, 18)
(2, 2, 16, 16, 36)	(3, 3, 4, 12, 50)	(5, 5, 10, 26, 26)	(12, 12, 12, 12, 24)

Table 2: Full 5-variable designs in 72 from full 4-variable designs in 36

Design in 24	Order 3 Circulant Matrices Used	Design in 72
1, 1, 1, 1, 5, 5, 10	a b-b; -c b b; c b b; -b b b; e d d; -e d d; -d d d;	1, 2, 9, 10, 50
1, 1, 1, 1, 5, 5, 10	a b-b; -c b b; -b c c; -c c c; e d d; -e d d; -d d d;	1, 5, 6, 10, 50
1, 1, 1, 1, 5, 5, 10	a b b; -a b b; c b-b; c b-b; e d d; -e d d; -d d d;	2, 2, 8, 10, 50
1, 1, 1, 1, 5, 5, 10	a b b; -a b b; -b b b; -b b b; d d d; e d-d; e d-d;	2, 10, 15, 45
1, 1, 1, 1, 5, 5, 10	a b b; -a b b; c b-b; c b-b; b b b; c b-b; -b b b;	2, 7, 63
1, 1, 1, 1, 5, 5, 10	a b b; -a b b; c b-b; c b-b; b b b; c b-b; c b-b;	2, 17, 53
1, 1, 1, 1, 3, 9, 8	b a a; -b a a; a a a; a a a; d d d; e d-d; -a a a;	2, 9, 27, 34
3, 3, 3, 3, 3, 3, 6	a b-b; -c b b; c b b; -b b b; d e e; d-e-e; -e e e;	3, 6, 6, 27, 30
3, 3, 3, 3, 3, 3, 6	a b-b; -c b b; b c c; -c c c; d e e; d-e-e; -e e e;	3, 6, 15, 18, 30
3, 3, 3, 3, 3, 3, 3, 3	a b-b; -c b b; c b b; -b b b; d e e; e-d-d; -e e e; -d d d;	3, 6, 18, 18, 27
3, 3, 3, 3, 3, 3, 3, 3	a b-b; -c b b; b-c-c; -c c c; d e e; e-d-d; -e e e; -d d d;	3, 15, 18, 18, 18
1, 1, 1, 1, 5, 5, 10	a b b; b-a-a; -a a a; -b b b; e d-d; d d d; -d d d;	5, 6, 6, 55
3, 3, 3, 3, 3, 3, 3, 3	a b b; -a b b; c b b; -c b b; d-b b; d-b b; d-b b; -b b b;	3, 6, 6, 9, 48

Table 3: Full orthogonal designs in order 72 constructed from full designs in order 24

(1, 2, 9, 60)	(2, 2, 18, 50)	(3, 6, 27, 36)	(5, 6, 10, 51)	(6, 9, 27, 30)
(1, 2, 10, 59)	(2, 8, 10, 52)	(3, 6, 30, 33)	(5, 6, 11, 50)	(6, 12, 24, 30)
(1, 2, 19, 50)	(2, 8, 12, 50)	(3, 9, 10, 50)	(5, 7, 10, 50)	(6, 15, 18, 33)
(1, 5, 6, 60)	(2, 9, 10, 51)	(3, 9, 12, 48)	(6, 6, 6, 54)	(6, 15, 21, 30)
(1, 5, 10, 56)	(2, 9, 11, 50)	(3, 12, 27, 30)	(6, 6, 9, 51)	(6, 18, 18, 30)
(1, 5, 16, 50)	(2, 9, 27, 34)	(3, 15, 18, 36)	(6, 6, 10, 50)	(6, 18, 21, 27)
(1, 6, 10, 55)	(2, 10, 10, 50)	(3, 15, 24, 30)	(6, 6, 12, 48)	(6, 18, 24, 24)
(1, 6, 15, 50)	(2, 10, 15, 45)	(3, 18, 18, 33)	(6, 6, 18, 42)	(9, 15, 18, 30)
(1, 9, 10, 52)	(3, 6, 6, 57)	(3, 18, 21, 30)	(6, 6, 24, 36)	(9, 18, 18, 27)
(1, 9, 12, 50)	(3, 6, 9, 54)	(3, 18, 24, 27)	(6, 6, 27, 33)	(12, 18, 18, 24)
(1, 10, 11, 50)	(3, 6, 15, 48)	(4, 8, 10, 50)	(6, 6, 30, 30)	(15, 18, 18, 21)
(2, 2, 8, 60)	(3, 6, 18, 45)	(5, 6, 6, 55)	(6, 9, 9, 48)	(18, 18, 18, 18)
(2, 2, 10, 58)				

Table 4: Full 4-variable $OD(72; s_1, s_2, s_3, 72 - s_1 - s_2 - s_3)$ constructed from full designs presented in table 3

Design in 12	Order 6 Matrices Used	Design in 72
2, 2, 4, 4	$xI + yC, xI - yC, zI + wC, zI - wC,$	4, 8, 20, 40
2, 2, 4, 4	$xI + yC, xI - yC, zI + wC, wI - zC,$	4, 20, 24, 24
2, 2, 4, 4	$xI + yC, yI - xC, zI + wC, zI - wC,$	8, 12, 12, 40

Table 5: Full orthogonal designs in order 72 constructed from full designs in order 12

Type	A_1 A_3 A_5 A_7	A_2 A_4 A_6 A_8	ZERO
(1,2,33,36)	(a, b, b, b, b, b, b, b) $(a, b, \bar{b}, \bar{b}, \bar{b}, \bar{b}, \bar{b}, \bar{b})$ $(e, e, \bar{e}, \bar{e}, \bar{e}, \bar{e}, e, e)$ $(e, \bar{e}, e, e, e, e, \bar{e}, \bar{e})$	$(e, e, \bar{e}, \bar{e}, \bar{e}, e, e, \bar{e})$ $(e, \bar{e}, e, e, e, \bar{e}, \bar{e}, e)$ $(b, b, \bar{b}, \bar{b}, \bar{b}, b, b, \bar{b})$ $(b, \bar{b}, b, b, c, \bar{b}, \bar{b}, b)$	NPAF $n = 9$
(1,9,26,36)	(a, b, a, b, b, b, b, b) $(e, \bar{e}, e, e, \bar{e}, \bar{e}, e, e)$ $(e, e, e, \bar{e}, e, e, \bar{e}, \bar{e})$ $(e, e, e, \bar{e}, e, \bar{e}, e, e)$	$(e, \bar{e}, e, e, \bar{e}, e, \bar{e}, \bar{e})$ $(b, \bar{b}, b, b, \bar{a}, a, a, a)$ $(b, a, b, \bar{a}, a, \bar{b}, b, b)$ $(b, b, b, \bar{b}, c, b, \bar{b}, \bar{b})$	NPAF $n = 9$
(2,2,2,2,16,16,16,16)	$(a, c, c, c, \bar{c}, d, d, d)$ $(f, h, h, h, \bar{h}, \bar{g}, \bar{g}, g, \bar{g})$ $(a, \bar{c}, \bar{c}, \bar{c}, c, \bar{d}, \bar{d}, d)$ $(f, \bar{h}, \bar{h}, \bar{h}, h, g, g, \bar{g})$	$(e, g, g, g, \bar{g}, h, h, h)$ $(b, d, d, d, \bar{c}, \bar{c}, c, \bar{c})$ $(e, \bar{g}, \bar{g}, \bar{g}, g, \bar{h}, \bar{h}, h)$ $(b, \bar{d}, \bar{d}, \bar{d}, d, c, c, \bar{c})$	NPAF $n = 9$
(3,3,6,6,27,27)	$(a, a, \bar{c}, d, d, d, d, d)$ $(a, \bar{d}, c, \bar{d}, \bar{d}, d, d, \bar{d})$ $(a, \bar{a}, d, \bar{d}, d, d, d, \bar{d})$ $(e, g, h, \bar{h}, h, h, h, \bar{h})$	$(e, \bar{e}, h, h, h, h, h, h)$ $(e, \bar{h}, g, \bar{h}, \bar{h}, h, h, \bar{h})$ $(e, e, \bar{g}, \bar{h}, \bar{h}, \bar{h}, h, h)$ $(a, c, d, \bar{d}, d, d, d, \bar{d})$	PAF $n = 9$
(3,3,3,3,30,30)	$(c, \bar{c}, \bar{c}, a, \bar{c}, \bar{b}, \bar{c}, \bar{c})$ $(c, \bar{c}, \bar{c}, \bar{c}, \bar{b}, \bar{c}, c, \bar{c})$ $(c, c, c, \bar{c}, a, c, c, \bar{c})$ $(c, \bar{c}, c, a, c, \bar{b}, \bar{c}, \bar{c})$	$(g, g, g, \bar{g}, e, g, g, \bar{g})$ $(g, \bar{g}, g, e, g, \bar{f}, \bar{g}, \bar{g})$ $(g, \bar{g}, \bar{g}, e, \bar{g}, \bar{f}, \bar{g}, g)$ $(g, \bar{g}, \bar{g}, \bar{g}, \bar{f}, \bar{g}, g, \bar{g})$	PAF $n = 9$
(3,3,15,51)	(d, d, d, a, b, b, d, d) $(d, \bar{d}, d, a, b, \bar{b}, \bar{d}, \bar{d})$ $(d, d, d, \bar{b}, a, b, d, \bar{d})$ $(d, \bar{d}, \bar{d}, \bar{b}, \bar{b}, \bar{d}, d, \bar{d})$	(d, d, d, b, d, b, d, d) $(d, \bar{d}, \bar{d}, \bar{b}, \bar{c}, \bar{b}, d, \bar{d})$ $(d, \bar{d}, \bar{d}, d, \bar{b}, c, \bar{d}, \bar{d})$ $(d, \bar{d}, d, d, b, \bar{c}, \bar{d}, \bar{d})$	PAF $n = 9$
(6,6,6,6,12,12,12,12)	$(c, d, a, c, d, b, \bar{c}, d, a)$ $(c, d, a, \bar{c}, d, \bar{a}, c, d, \bar{a})$ $(c, \bar{d}, b, c, \bar{d}, \bar{a}, \bar{c}, d, b)$ $(c, \bar{d}, b, c, d, \bar{b}, c, \bar{d}, \bar{b})$	$(g, h, e, g, h, f, \bar{g}, h, e)$ $(g, h, e, \bar{g}, h, \bar{e}, g, h, \bar{e})$ $(g, \bar{h}, f, g, \bar{h}, \bar{e}, \bar{g}, h, f)$ $(g, \bar{h}, f, g, h, \bar{f}, g, \bar{h}, \bar{f})$	NPAF $n = 9$
(9,9,9,9,18,18)	$(a, a, a, \bar{a}, b, b, b, b, c)$ $(g, g, g, \bar{g}, e, e, \bar{e}, e, \bar{e})$ $(f, f, f, \bar{f}, \bar{e}, \bar{e}, e, \bar{e}, \bar{e})$ $(a, a, a, \bar{a}, \bar{c}, \bar{c}, c, \bar{c}, b)$	$(e, e, e, \bar{e}, \bar{g}, \bar{g}, g, \bar{g}, f)$ $(c, c, c, \bar{c}, a, a, \bar{a}, a, \bar{a})$ $(b, b, b, \bar{b}, \bar{a}, \bar{a}, a, \bar{a}, \bar{a})$ $(e, e, e, \bar{e}, f, f, \bar{f}, f, g)$	NPAF $n = 9$

Table 6: New full orthogonal designs in order 72 constructed from new amicable sets of eight matrices

(1, 2, 33, 36)	(2, 16, 16, 38)	(3, 6, 9, 54)	(6, 6, 12, 48)	(8, 16, 16, 32)
(1, 9, 26, 36)	(2, 16, 18, 36)	(3, 6, 27, 36)	(6, 6, 18, 42)	(9, 9, 9, 45)
(2, 2, 2, 66)	(2, 16, 20, 34)	(3, 6, 30, 33)	(6, 6, 24, 36)	(9, 9, 18, 36)
(2, 2, 4, 64)	(2, 16, 22, 32)	(3, 9, 27, 33)	(6, 6, 27, 33)	(9, 9, 27, 27)
(2, 2, 16, 52)	(2, 18, 18, 34)	(3, 9, 30, 30)	(6, 6, 30, 30)	(9, 18, 18, 27)
(2, 2, 18, 50)	(2, 18, 20, 32)	(3, 12, 27, 30)	(6, 9, 27, 30)	(12, 12, 12, 36)
(2, 2, 20, 48)	(3, 3, 3, 63)	(3, 15, 27, 27)	(6, 12, 12, 42)	(12, 12, 18, 30)
(2, 2, 32, 36)	(3, 3, 6, 60)	(4, 4, 16, 48)	(6, 12, 18, 36)	(12, 12, 24, 24)
(2, 2, 34, 34)	(3, 3, 12, 54)	(4, 4, 32, 32)	(6, 12, 24, 30)	(12, 18, 18, 24)
(2, 4, 16, 50)	(3, 3, 15, 51)	(4, 16, 16, 36)	(6, 12, 27, 27)	(16, 16, 16, 24)
(2, 4, 18, 48)	(3, 3, 27, 39)	(4, 16, 18, 34)	(6, 16, 16, 34)	(16, 16, 18, 22)
(2, 4, 32, 34)	(3, 3, 30, 36)	(4, 16, 20, 32)	(6, 16, 18, 32)	(16, 16, 20, 20)
(2, 6, 16, 48)	(3, 3, 33, 33)	(4, 18, 18, 32)	(6, 18, 18, 30)	(16, 18, 18, 20)
(2, 6, 32, 32)	(3, 6, 6, 57)	(6, 6, 6, 54)	(6, 18, 24, 24)	(18, 18, 18, 18)

Table 7: Full 4-variable $OD(72; s_1, s_2, s_3, 72 - s_1 - s_2 - s_3)$ constructed from full designs presented in table 6

s_1, s_2, s_3	s_1, s_2, s_3	s_1, s_2, s_3	s_1, s_2, s_3	s_1, s_2, s_3	s_1, s_2, s_3
1, 8, 63	5, 14, 53	7, 24, 41	9, 19, 44	12, 22, 38	17, 24, 31
1, 14, 57	5, 18, 49	7, 25, 40	9, 20, 43	13, 14, 45	17, 25, 30
1, 24, 47	5, 20, 47	7, 26, 39	9, 22, 41	13, 19, 40	17, 27, 28
1, 25, 46	5, 24, 43	7, 27, 38	9, 23, 40	13, 21, 38	18, 23, 31
1, 30, 41	5, 25, 42	7, 28, 37	9, 24, 39	13, 22, 37	19, 20, 33
1, 31, 40	5, 27, 40	7, 30, 35	9, 31, 32	13, 24, 35	19, 21, 32
1, 32, 39	5, 28, 39	7, 31, 34	10, 19, 43	13, 25, 34	19, 22, 31
2, 23, 47	5, 29, 38	7, 32, 33	10, 29, 33	13, 27, 32	19, 23, 30
3, 8, 61	5, 30, 37	8, 9, 55	11, 12, 49	14, 15, 43	19, 24, 29
3, 11, 58	5, 32, 35	8, 11, 53	11, 14, 47	14, 17, 41	19, 25, 28
3, 13, 56	5, 33, 34	8, 13, 51	11, 18, 43	14, 19, 39	19, 26, 27
3, 14, 55	6, 7, 59	8, 15, 49	11, 19, 42	14, 20, 38	20, 21, 31
3, 17, 52	6, 13, 53	8, 17, 47	11, 21, 40	14, 25, 33	21, 22, 29
3, 26, 43	6, 19, 47	8, 19, 45	11, 22, 39	14, 28, 30	23, 24, 25
3, 28, 41	7, 8, 57	8, 21, 43	11, 23, 38	15, 17, 40	
3, 29, 40	7, 9, 56	8, 23, 41	11, 24, 37	15, 19, 38	
3, 31, 38	7, 13, 52	8, 25, 39	11, 26, 35	15, 23, 34	
3, 32, 37	7, 14, 51	8, 27, 37	11, 28, 33	15, 25, 32	
4, 5, 63	7, 18, 47	8, 29, 35	11, 29, 32	15, 28, 29	
4, 19, 49	7, 20, 45	8, 31, 33	12, 13, 47	17, 20, 35	
5, 8, 59	7, 21, 44	9, 14, 49	12, 17, 43	17, 22, 33	
5, 9, 58	7, 22, 43	9, 17, 46	12, 19, 41	17, 23, 32	

Table 8: The existence of these 124 full $OD(72; s_1, s_2, 72 - s_1 - s_2)$ is not yet established.

4 Summary

We have found new designs in order 72 and shown that of 2700 possible $OD(72; s_1, s_2, s_3, 72 - s_1 - s_2 - s_3)$ 335 are known: of 432 possible $OD(72; s_1, s_2, 72 - s_1 - s_2)$ 308 are known; and all possible $OD(72; s_1, 72 - s_1)$ are known.

References

- [1] A.V.Geramita, J.M.Geramita, and J.Seberry Wallis, Orthogonal designs, *Linear and Multilinear Algebra*, 3 (1976), 281-306.
- [2] A.V.Geramita, and J.Seberry, *Orthogonal Designs: Quadratic Forms and Hadamard Matrices*, Marcel Dekker, New York-Basel, 1979.
- [3] S. Georgiou, C. Koukouvinos, M. Mitrouli and J. Seberry, Necessary and sufficient conditions for three and four variable orthogonal designs in order 36, *J. Statist. Plann. Inference*, (to appear).
- [4] H. Kharaghani, Arrays for orthogonal designs, *J. Combin. Designs*, to appear.
- [5] C.Koukouvinos, M.Mitrouli, J.Seberry, and P.Karabelas, On sufficient conditions for some orthogonal designs and sequences with zero autocorrelation function, *Australas. J. Combin.*, 13 (1996), 197-216.
- [6] C. Koukouvinos, N. Platis and Jennifer Seberry, Necessary and sufficient conditions for some two variable orthogonal designs in order 36, *Congr. Numerantium*, 114 (1996), 129-139.
- [7] J. Seberry Wallis, On the existence of Hadamard matrices, *J. Combin. Theory Ser. A*, 21 (1976), 188-195.