

Orthogonal Designs from Negacyclic Matrices

Ken Finlayson and Jennifer Seberry
School of IT and Computer Science
University of Wollongong
NSW 2522
Australia

19 March 2003

Abstract

We study the use of negacyclic matrices to form orthogonal designs and hence Hadamard matrices. We give results for all possible tuple for order 12, all but 3 for order 20 and all but 3 for order 28.

Key words and phrases: Orthogonal designs, negacyclic matrices, Hadamard matrices, variant Goethals-Seidel array

AMS Subject Classification:

1 Introduction

An *orthogonal design* of order n and type (s_1, s_2, \dots, s_k) in variables x_1, x_2, \dots, x_k , denoted $OD(n; s_1, s_2, \dots, s_k)$, is a matrix A of order n with entries in the set $\{0, \pm x_1, \pm x_2, \dots, \pm x_k\}$ satisfying

$$AA^T = \left(\sum_{i=1}^k s_i x_i^2\right) I_n,$$

where I_n is the identity matrix of order n . Alternatively, the rows of A are formally orthogonal and each row has precisely s_i entries of the type $\pm x_i$. In [1], where this was first defined, it was mentioned that

$$A^T A = \left(\sum_{i=1}^k s_i x_i^2\right) I_n$$

and so our alternative description of A applies equally well to the columns of A .

An *Hadamard matrix* H of order n is a square $(1, -1)$ matrix having inner product of distinct rows zero. Hence $HH^T = nI_n$. We note that $n = 1, 2$ or $n \equiv 0 \pmod{4}$.

A matrix $A \pm I$ is *skew-type* if A has zero diagonal and $A^T = -A$. A skew-type Hadamard matrix is said to be *skew-Hadamard*.

Circulant matrices of order n are polynomials in the shift matrix

$$S = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & & 0 \\ \vdots & & & & \vdots \\ 0 & 0 & 0 & & 1 \\ 1 & 0 & 0 & & 0 \end{pmatrix}.$$

Negacyclic matrices of order n are polynomials in the negashift matrix

$$N = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & & 0 \\ \vdots & & & & \vdots \\ 0 & 0 & 0 & & 1 \\ -1 & 0 & 0 & & 0 \end{pmatrix}.$$

The *back-diagonal matrix* R of order n is the matrix whose elements r_{ij} are given by

$$r_{ij} = \begin{cases} 1 & \text{if } i + j = n + 1, \\ 0 & \text{otherwise} \end{cases}$$

where $i, j = 1, \dots, n$. We note that if A, B are polynomial in S then $A(BR)^T = (BR)A^T$. We show below that the same is true if A, B are polynomial in N .

Lastly, we define the *nonperiodic autocorrelation function*. Let

$$X = \{\{a_{11}, \dots, a_{1n}\}, \{a_{21}, \dots, a_{2n}\}, \dots, \{a_{m1}, \dots, a_{mn}\}\}$$

be m sequences of commuting variables of length n . Then the nonperiodic autocorrelation function of the family of sequences X (denoted N_X) is the function defined by

$$N_X(j) = \sum_{i=1}^{n-j} (a_{1,i}a_{1,i+j} + a_{2,i}a_{2,i+j} + \cdots + a_{m,i}a_{m,i+j}).$$

2 Preliminary results

We note some properties of the negashift matrix N given above:

$$(N^i)^T = -N^{n-i} \quad \text{and} \quad N^i R = -RN^{n-i}.$$

Hence we have:

Lemma 1 $N^i(N^j R)^T = (N^j R)(N^i)^T$.

Proof. $N^i(N^j R)^T = N^i R(N^j)^T = -N^i R N^{n-j} = N^i N^j R = N^j N^i R = -N^j R N^{n-i} = (N^j R)(N^i)^T$. \square

Lemma 2 Suppose A, B are polynomial in S or N then $A(BR)^T = (BR)A^T$.

Proof. The result for S , circulant, can be found in Wallis [6]. For N , negacyclic, we note A and B are polynomials in N so by repeated applications of Lemma 1 we have the result. \square

We now develop some properties of negacyclic matrices.

Theorem 1 Let R_i, R_j be two rows of a negacyclic matrix of dimension n , where $1 < i < j \leq n$. Then $R_i R_j^T = R_1 R_{1+j-i}^T$.

Proof. Let $i = 1 + s$ and $j = 1 + s + t$. If we write $R_1 = (x_1, \dots, x_n)$, we have

$$\begin{aligned} R_{1+s} &= (-x_{n-s+1}, \dots, -x_n, x_1, \dots, x_{n-s}) \\ R_{1+t} &= (-x_{n-t+1}, \dots, -x_n, x_1, \dots, x_{n-t}) \\ R_{1+s+t} &= (-x_{n-s-t+1}, \dots, -x_n, x_1, \dots, x_{n-s-t}) \end{aligned}$$

We note that

$$\begin{aligned} R_{1+s} &= \overbrace{(-x_{n-s}, \dots, -x_n)}^s, \overbrace{(x_1, \dots, x_t, x_{t+1}, \dots, x_{n-s})}^t \\ R_{1+s+t} &= \overbrace{(-x_{n-s-t+1}, \dots, -x_{n-t})}^s, \overbrace{(-x_{n-t+1}, \dots, -x_n, x_1, \dots, x_{n-s-t})}^t \end{aligned}$$

Then

$$\begin{aligned} R_i R_j^T &= R_{1+s} R_{1+s+t}^T \\ &= x_{n-s+1} x_{n-s-t+1} + \dots + x_n x_{n-t} - x_1 x_{n-t+1} - \dots - x_t x_n \\ &\quad + x_{t+1} x_1 + \dots + x_{n-s} x_{n-s-t} \\ &= -x_1 x_{n-t+1} - \dots - x_t x_n + x_{t+1} x_1 + \dots + x_{n-s} x_{n-s-t} \\ &\quad + x_{n-s+1} x_{n-s-t+1} + \dots + x_n x_{n-t} \\ &= -x_1 x_{n-t+1} - \dots - x_t x_n + x_{t+1} x_1 + \dots + x_n x_{n-t} \\ &= R_1 R_{1+t}^T \\ &= R_1 R_{1+j-i}^T \end{aligned}$$

□

Corollary 1 For a negacyclic matrix of odd dimension n , there are only n inner products of interest, i.e. each row with the first. More precisely, for a negacyclic matrix of odd dimension n , there are $\frac{n+1}{2}$ distinct inner products, $p_1, p_2, p_3, \dots, p_{\frac{n+1}{2}}$, where $p_i = R_1 R_i^T$. The remaining $\frac{n-1}{2}$ inner products are related by the property that $-p_{\frac{n+1}{2}+q} = p_{\frac{n+3}{2}-q}$, for $1 \leq q \leq \frac{n-1}{2}$.

Corollary 2 For a negacyclic matrix of odd dimension n , $R_1 R_i^T = -R_1 R_{n-i+2}^T$ for $1 < i \leq n$.

Definition 1 Let L, M be two negacyclic matrices of dimension n . We say L and M are in the same inner product equivalence class if for every $1 \leq i, j \leq n$, $R_i R_j^T$ of L equals $R_i R_j^T$ of M .

Theorem 2 Let M be a negacyclic matrix of dimension n , with first row

$$m_1, m_2, \dots, m_n.$$

Then the negacyclic matrices with first rows

$$-m_1, -m_2, \dots, -m_n \quad \text{and} \quad m_n, m_{n-1}, \dots, m_1$$

(denoted $-M$ and M^* respectively) are in the same inner product equivalence class as M .

Proof. By Theorem 1, we need only consider the inner products of rows R_1, R_{1+j} for $1 \leq j \leq n-1$. We observe that for the matrix M ,

$$\begin{aligned} R_1 &= m_1 & m_2 & \cdots & m_j & m_{j+1} & m_{j+2} & \cdots & m_n \\ R_{1+j} &= -m_{n-j+1} & -m_{n-j+2} & \cdots & -m_n & m_1 & m_2 & \cdots & m_{n-j}. \end{aligned}$$

Likewise, for the matrix $-M$,

$$\begin{aligned} R_1 &= -m_1 & -m_2 & \cdots & -m_j & -m_{j+1} & -m_{j+2} & \cdots & -m_n \\ R_{1+j} &= m_{n-j+1} & m_{n-j+2} & \cdots & m_n & -m_1 & -m_2 & \cdots & -m_{n-j}. \end{aligned}$$

Similarly, for the matrix M^* ,

$$\begin{aligned} R_1 &= m_n & m_{n-1} & \cdots & m_{n-j+1} & m_{n-j} & m_{n-j-1} & \cdots & m_1 \\ R_{1+j} &= -m_j & -m_{j-1} & \cdots & -m_1 & m_n & m_{n-1} & \cdots & m_{j+1}. \end{aligned}$$

The inner product of R_1 with R_{1+j} is the same in all three cases, namely

$$\begin{aligned} R_1 R_{1+j}^T &= -m_1 m_{n-j+1} - m_2 m_{n-j+2} - \cdots - m_j m_n \\ &\quad + m_1 m_{j+1} + m_2 m_{j+2} + \cdots + m_{n-j} m_n \\ &= -\sum_{i=1}^j m_i m_{n-j+i} + \sum_{i=1}^{n-j} m_i m_{i+j}. \end{aligned}$$

Hence $M, -M$ and M^* are in the same inner product equivalence class. \square

Corollary 3 *Given a negacyclic matrix M , the negacyclic matrix $-M^*$ is in the same inner product equivalence class as M .*

Theorem 3 *Let M be a negacyclic matrix with first row $R_1 = (m_1, \dots, m_n)$. Then every negacyclic matrix M' with a first row R'_1 equal to a negashift of R_1 is in the same inner product equivalence class as M . (There are $2n-1$ negashifts of R_1 .)*

Proof. The $2n-1$ negashifts of R_1 are

$$R_2, \dots, R_n, -R_1, \dots, -R_n$$

where R_i is the i th row of the matrix M .

By Theorem 2, the negacyclic matrix with first row equal to $-R_1$ is in the same inner product equivalence class as M . Likewise, the negacyclic matrices with first rows equal to $-R_2, \dots, -R_n$ are in the same inner product equivalence class as the negacyclic matrices with first rows R_2, \dots, R_n respectively. Thus we need consider only the negacyclic matrices with first rows $R'_1 = R_i$, where $2 \leq i \leq n$.

Let M' be a negacyclic matrix with first row $R'_1 = R_i$ for some i in the range $2 \leq i \leq n$. By Corollary 1, M' has only $\frac{n+1}{2}$ distinct inner products, namely $R'_1 (R'_j)^T$, for $2 \leq j \leq \frac{n+1}{2}$.

Now, $R'_1 = R_i$, and

$$R'_{\frac{n+1}{2}} = \begin{cases} R_{i+\frac{n-1}{2}} & \text{for } 2 \leq i \leq \frac{n+1}{2} \\ -R_{i+\frac{n-1}{2}-n} & \text{for } \frac{n+3}{2} \leq i \leq n. \end{cases}$$

We consider the two cases in turn.

Case I: When $2 \leq i \leq \frac{n+1}{2}$, the rows $R'_1, \dots, R'_{\frac{n+1}{2}}$ are all from M . Then the inner product $R'_1(R'_j)^T$ for $2 \leq j \leq \frac{n+1}{2}$ is:

$$\begin{aligned} R'_1(R'_j)^T &= R_i R_{i+j-1}^T \\ &= R_1 R_j^T \quad (\text{by Theorem 1}). \end{aligned}$$

Thus the negacyclic matrices M' with first row $R'_1 = R_i$ for $2 \leq i \leq \frac{n+1}{2}$ are in the same inner product equivalence class as M .

Case II: When $\frac{n+3}{2} \leq i \leq n$, the rows R'_1, \dots, R'_{n-i+1} are from M , and the rows $R'_{n-i+2}, \dots, R'_{\frac{n+1}{2}}$ are from $-M$. For $2 \leq j \leq n-i+1$, $R'_1(R'_j)^T = R_1 R_j^T$ by the argument presented in Case I. For $n-i+2 \leq j \leq \frac{n+1}{2}$, we have

$$\begin{aligned} R'_1(R'_j)^T &= R_i R_{i+j-1-n}^T \\ &= -R_1 R_{n-j+2} \quad (\text{observe } i+j-1-n < i \\ &\quad \text{and then apply Theorem 1}) \\ &= R_1 R_j \quad (\text{by Corollary 2}). \end{aligned}$$

Thus the negacyclic matrices M' with first row $R'_1 = R_i$ for $\frac{n+3}{2} \leq i \leq n$ are in the same inner product equivalence class as M .

Hence the $2n-1$ negacyclic matrices which have a first row equal to a negashift of the first row of M are all in the same inner product equivalence class as M . \square

We note from Wallis and Whiteman [7] that circulant can be replaced by group-type or type 1 in abelian groups so that all results that follow for circulant also follow for group-type or type 1. Similarly we observe that group-type or type 1 negacyclic can be used instead of negacyclic and the corresponding results hold. So we have, modifying Goethals-Seidel construction [6]:

Theorem 4 *Suppose there exist four negacyclic $(1, -1)$ matrices A, B, C, D of order n . Further, suppose*

$$AA^T + BB^T + CC^T + DD^T = 4nI_n.$$

Then

$$SF = \begin{bmatrix} A & BR & CR & DR \\ -BR & A & D^T R & -C^T R \\ -CR & -D^T R & A & B^T R \\ -DR & C^T R & -B^T R & A \end{bmatrix} \quad (1)$$

is an Hadamard matrix of order $4n$ of SF type. (Here R is the back diagonal matrix.) If A is of skew-type, then SF is skew-Hadamard.

3 Our results

Definition 2 *A set of four negacyclic matrices A, B, C, D is said to be suitable if*

$$AA^T + BB^T + CC^T + DD^T = fI$$

for some f .

Theorem 5 *If there exist four sequences with zero non-periodic autocorrelation function, then there exist four suitable negacyclic matrices.*

Proof. Let there be four sequences of length n , denoted

$$X = \{\{a_1, a_2, \dots, a_n\}, \{b_1, b_2, \dots, b_n\}, \{c_1, c_2, \dots, c_n\}, \{d_1, d_2, \dots, d_n\}\},$$

with zero non-periodic autocorrelation function. We now treat each of these sequences as the first row of a negacyclic matrix. The matrices generated in this way are denoted A, B, C, D respectively. We consider the sum

$$AA^T + BB^T + CC^T + DD^T.$$

For a negacyclic matrix M , the element (i, j) of the product MM^T is equal to $R_i R_j^T$, where R_i, R_j are the i th and j th rows of M respectively. By Theorem 1, it is sufficient to consider only the products $R_1 R_{1+j}^T$, for $1 \leq j \leq n-1$. We recall that

$$R_1 R_{1+j}^T = - \sum_{i=1}^j m_i m_{n-j+i} + \sum_{i=1}^{n-j} m_i m_{i+j}$$

(as shown in the proof of Theorem 2).

Thus the sum of the products $R_1 R_{1+j}^T$ of A, B, C and D , for $1 \leq j \leq n-1$, is equal to

$$\begin{aligned} & - \sum_{i=1}^j (a_i a_{n-j+i} + b_i b_{n-j+i} + c_i c_{n-j+i} + d_i d_{n-j+i}) \\ & + \sum_{i=1}^{n-j} (a_i a_{i+j} + b_i b_{i+j} + c_i c_{i+j} + d_i d_{i+j}) \\ & = -N_X(n-j) + N_X(j) \\ & = 0 \end{aligned}$$

By Theorem 1, the off-diagonal element (i, j) , where $i \neq j$, of the sum $AA^T + BB^T + CC^T + DD^T$, is equal to element $(1, 1+j-i)$. As has just been shown, these elements all equal to 0. Thus all off-diagonal elements of the sum are 0.

Lastly, we consider the diagonal elements. It is clear that element $(j, j) = \sum_{i=1}^n (a_i^2 + b_i^2 + c_i^2 + d_i^2)$ for $1 \leq j \leq n$. Write this sum as f . Thus we have

$$AA^T + BB^T + CC^T + DD^T = fI$$

and so the negacyclic matrices A, B, C, D are suitable. \square

Corollary 4 *The four suitable negacyclic matrices A, B, C, D that were constructed in the last proof satisfy*

$$AA^T + BB^T + CC^T + DD^T = 4nI_n.$$

Hence, by Theorem 4, we can produce an Hadamard matrix of order $4n$ of SF-type.

We have found that suitable negacyclic matrices are not limited to those generated from sequences with a zero non-periodic autocorrelation function. The appendix contains a listing of negacyclic sequences of order $n = 3, 5, 7$ which are suitable, and so yield SF-type Hadamard matrices and orthogonal designs of order $4n$.

References

- [1] A. V. Geramita, J. M. Geramita and J. Seberry. Orthogonal designs. *Linear and Multi-linear algebra*. **3**: 281–306, 1975/76.
- [2] A. V. Geramita and J. Seberry. *Orthogonal designs: Quadratic forms and Hadamard matrices*. New York-Basel, 1979.
- [3] A. V. Geramita and J. H. Verner. Orthogonal designs with zero diagonal. *Canad. J. Math.* **28**: 215–225, 1976.
- [4] J. Seberry. On skew Hadamard matrices. *Ars Combinatoria*. **6**: 255–275, 1978.
- [5] J. Seberry and R. Craigen. Orthogonal designs. In *CRC Handbook of Combinatorial Designs*, C. J. Colbourn and J. H. Dinitz (eds), CRC Press, 1996, 400–406.
- [6] J. S. Wallis. Hadamard matrices. In *Combinatorics: Room Squares, Sum-Free Sets and Hadamard Matrices*. W. D. Wallis, A. P. Street and J. S. Wallis, Lecture Notes in Computer Science, **292**, Springer-Verlag, 1972.
- [7] J. S. Wallis and A. L. Whiteman. Some classes of Hadamard matrices with constant diagonal. *Bulletin of the Australian Mathematical Society*. **7**: 233–249, 1972.

A Negacyclic sequences with zero autocorrelation function

Order 12

Design	A	B	C	D
(1,1,1,1)	a 0 0	b 0 0	c 0 0	d 0 0
(1,1,1,4)	a 0 0	b 0 0	c d d	0 d -d
(1,1,1,9)	a d d	b d d	c d d	d -d d
(1,1,2,2)	a 0 0	b 0 0	c -d 0	c d 0
(1,1,2,8)	a d d	b d d	-c -d d	c -d d
(1,1,4,4)	a d d	b c c	0 -c c	0 -d d
(1,1,5,5)	a d d	b c c	d -c c	-c -d d
(1,2,2,4)	0 d -d	a d d	b c 0	b -c 0
(1,2,3,6)	a d -d	c d d	c a -d	c -a d
(2,2,2,2)	a b 0	a -b 0	c d 0	c -d 0
(2,2,4,4)	a b b	d b -b	c d d	-b d -d
(3,3,3,3)	a b c	b -c d	c d -a	d a -b

Order 20

Design	A, C	B, D
(1,1,1,1)	a 0 0 0 0	b 0 0 0 0
	c 0 0 0 0	d 0 0 0 0
(1,1,1,4)	a 0 0 0 0	b 0 0 0 0
	c 0 d d 0	0 0 d -d 0
(1,1,1,9)	a 0 d d 0	b 0 d d 0
	c d 0 0 d	d -d d 0 0
(1,1,2,2)	a 0 0 0 0	b 0 0 0 0
	c -d 0 0 0	d c 0 0 0

(1,1,2,8)	a 0 d d 0	b 0 d d 0
	c 0 d -d 0	-c 0 d -d 0
(1,1,4,4)	a 0 c c 0	0 0 c -c 0
	b 0 d d 0	0 0 d -d 0
(1,1,4,9)	a 0 d d 0	b 0 d d 0
	0 -d c -c -d	d -d c c d
(1,1,5,5)	a 0 c c 0	b 0 d d 0
	c 0 d -d 0	-d 0 c -c 0
(1,1,8,8)	a -c d d -c	b -c -d -d -c
	0 -c d -d c	0 -c -d d c
(1,1,9,9)	a -c d d -c	b -d -c -c -d
	-c -c c -c c	-d -d d -d d
(1,2,2,4)	a 0 d d 0	0 0 d -d 0
	b -c 0 0 0	c b 0 0 0
(1,2,2,9)	a -d 0 0 -d	d -d d 0 0
	b -c d d 0	c b 0 -d -d
(1,2,3,6)	a 0 d d 0	b -c 0 -d 0
	b c 0 d 0	c 0 -d d 0
(1,2,4,8)	a 0 d d 0	b -c d -d -c
	b 0 -d d 0	d d c 0 c
(1,2,8,9)	a -d c c -d	d -d d -c -c
	c -b d d c	b c -c -d -d
(1,4,4,4)	a 0 b b 0	0 0 b -b 0
	0 -c d d c	0 -c d -d -c
(1,4,5,5)	a -d 0 0 -d	-d -c 0 0 c
	c -d b b d	0 -c b -b -c
(1,5,5,9)	a -b c c -b	-c -b d d b
	b -d c -c -d	-d -d d -d d
(2,2,2,2)	a -b 0 0 0	b a 0 0 0
	c -d 0 0 0	d c 0 0 0
(2,2,2,8)	a -d b d 0	a d -b -d 0
	c 0 d 0 d	c 0 -d 0 -d
(2,2,4,4)	a -b 0 0 0	b a 0 0 0
	0 -c d d c	0 -c d -d -c
(2,2,4,9)	a -b d d 0	a b d d 0
	0 -d c -c -d	d -d c c d
(2,2,8,8)	a -d c -d -c	a d -c d c
	b -c -d -c d	b c d c -d
(2,3,4,6)	b -d a 0 0	b -d -a 0 0
	-b -d c c d	0 -d c -c -d
(2,3,6,9)	b -d c -a -d	a -d -d b -c
	-b -d -d -c c	d -d -c -c d
(2,4,4,8)	c 0 b -d d	c 0 b d -d
	b -a -c -d -d	b a -c d d
(2,5,5,8)	c -b d d b	a -c d -d -c
	-a -b d -d -b	-b -c d d c
(3,3,3,3)	a -b c 0 0	a b 0 -d 0
	a 0 -c d 0	b c d 0 0
(3,3,6,6)	0 -d -a -c -b	0 -d a c -b
	b -c d -d -c	a -c d d c
(4,4,4,4)	0 -a b b a	0 -a b -b -a
	0 -c d d c	0 -c d -d -c
(4,4,5,5)	d -c a a c	0 -d a -a -d
	0 -c b -b -c	-c -d b b d
(5,5,5,5)	a -b b -d -d	-b -a a -c -c
	d -c c a a	-c -d d b b
(1,1,13)	a -c -c -c -c	0 -c -c c -c
	b -c 0 0 -c	0 -c c -c 0
(1,2,17)	a -c -c -c -c	b -c -c c -c
	b c c -c c	c c -c c -c

(1,2,11)	a 0 c c 0	b 0 c c -c
	b 0 -c -c c	0 -c 0 -c c
(1,3,14)	a -c -c -c -c	-b -c c -c 0
	c -c c -b -c	c -c -c 0 b
(1,4,13)	a -c -c -c -c	0 -c -c c -c
	0 -c b -b -c	b -c c -c -b
(1,6,11)	a -c b b -c	c -c b -b -c
	c b c -c 0	c 0 c c b
(1,8,11)	a -c -b -b -c	-c -c b -b -c
	c b c -c b	c -b c c b
(2,5,7)	c -c -c 0 -a	a -c b b 0
	b -b 0 -c 0	c b 0 -c 0
(3,6,8)	0 -b c b -c	0 -b c -a c
	a -b -c 0 -c	a b c b -c
(7,10)	0 -a a b -b	b -a a a a
	0 -a a a b	b 0 b b a

Order 28

Design	A, C	B, D
(1,1,1,1)	a 0 0 0 0 0	b 0 0 0 0 0 0
	c 0 0 0 0 0	d 0 0 0 0 0 0
(1,1,1,4)	a 0 0 0 0 0	b 0 0 0 0 0 0
	d -c -d 0 0 0	d 0 d 0 0 0 0
(1,1,1,9)	d -a -d 0 0 0	d -b -d 0 0 0 0
	d 0 c 0 -d 0	d 0 d 0 d 0 0
(1,1,1,16)	a -d d d d -d	b 0 d 0 0 d 0
	c 0 d 0 0 d 0	0 -d d d d -d d
(1,1,1,25)	a -d d d d -d	b -d d d d -d
	c -d d d d -d	-d -d d -d d -d d
(1,1,2,2)	a 0 0 0 0 0	b 0 0 0 0 0 0
	c -d 0 0 0 0	c d 0 0 0 0 0
(1,1,2,8)	d -a -d 0 0 0	d -b -d 0 0 0 0
	d -c d 0 0 0	d c d 0 0 0 0
(1,1,2,18)	a -b b b b -b	c -b b b b -b
	d -b b 0 b 0	-d -b b 0 b 0 0
(1,1,4,4)	a 0 0 0 0 0	b 0 0 0 0 0 0
	c -c d d 0 0	d -d -c -c 0 0 0
(1,1,4,9)	a -d d d d -d	0 -d d 0 d 0 0
	b 0 0 -c -c 0	0 0 0 -c c 0 0
(1,1,4,16)	b 0 b 0 b 0 b	b 0 b -a -b 0 -b
	b -c -b 0 -b -c b	b -c -b -d b c -b
(1,1,5,5)	a 0 0 0 0 0	b 0 0 0 0 0 0
	c -c -c -d 0 -d	d -d -d c 0 c 0
(1,1,8,8)	c -d a d -c 0	-c -d b d c 0 0
	c -d 0 -d c 0	c d 0 d c 0 0
(1,1,9,9)	a -c c c c -c	0 -c c 0 c 0 0
	b -d d d d -d	0 -d d 0 d 0 0
(1,1,10,10)	d -a a a a -a	c -b b b b -b
	-a -b b 0 b 0	b -a a 0 a 0 0
(1,1,13,13)	-a -b b -a b -a a	-b a -a -b -a -b b
	c -a a a a -a	d -b b b b -b
(1,2,2,4)	b -a -b 0 0 0	b 0 b 0 0 0 0
	c -d 0 0 0 0	c d 0 0 0 0 0
(1,2,2,9)	a -d d d d -d	0 -d d 0 d 0 0
	b -c 0 0 0 0	b c 0 0 0 0 0
(1,2,2,16)	a -d d d d -d	b 0 d -c 0 d 0
	b 0 d c 0 d 0	0 -d d d d -d d
(1,2,3,6)	a -b c 0 0 0	a -b -c 0 0 0 0
	b a b 0 0 0	b -d -b 0 0 0 0

(1,2,4,8)	c -a -c 0 0 0 0	c 0 c 0 0 0 0
	d -b -d -d 0 -d 0	d -b -d d 0 d 0
(1,2,8,9)	a -d d d d d -d	0 -d d 0 d 0 0
	-b -b b -c b 0 0	-b -b b c b 0 0
(1,3,6,8)	-c -c c 0 c -a b	c c -c 0 -c -a b
	b a b 0 0 0 0	b -d -b 0 0 0 0
(1,4,4,4)	b -a -b 0 0 0 0	b 0 b 0 0 0 0
	c -c d d 0 0 0	d -d -c -c 0 0 0
(1,4,4,9)	a -d d d d d -d	0 -d d 0 d 0 0
	c -c b b 0 0 0	b -b -c -c 0 0 0
(1,4,5,5)	b -a -b 0 0 0 0	b 0 b 0 0 0 0
	c -c -c -d 0 -d 0	d -d -d c 0 c 0
(1,5,5,9)	a -d d d d d -d	0 -d d 0 d 0 0
	b -b -b -c 0 -c 0	c -c -c b 0 b 0
(2,2,2,2)	a -b 0 0 0 0 0	a b 0 0 0 0 0
	c -d 0 0 0 0 0	c d 0 0 0 0 0
(2,2,8,2)	a -b 0 0 0 0 0	a b 0 0 0 0 0
	c -d -c -c 0 -c 0	c -d -c c 0 c 0
(2,2,2,18)	a -d -d 0 b -d -d	a -d -d 0 -b -d -d
	c -d d -d d d 0	-c -d d -d d d 0
(2,2,4,4)	a -b 0 0 0 0 0	a b 0 0 0 0 0
	c -c d d 0 0 0	d -d -c -c 0 0 0
(2,2,4,16)	a -b -a -a c -a 0	a -b -a -a -c -a 0
	a -b -a a d a 0	a -b -a a -d a 0
(2,2,5,5)	a -b 0 0 0 0 0	a b 0 0 0 0 0
	c -c -c -d 0 -d 0	d -d -d c 0 c 0
(2,2,8,8)	a -b -a -a 0 -a 0	a -b -a a 0 a 0
	c -d -c -c 0 -c 0	c -d -c c 0 c 0
(2,2,9,9)	d -b -d -c a c 0	d -b -d c -a -c 0
	d -c 0 -c -d -c 0	-c -d 0 -d c -d 0
(2,2,10,10)	c -c -c -d a -d 0	c -c -c -d -a -d 0
	d -d -d c b c 0	d -d -d c -b c 0
(2,3,4,6)	a -d 0 d a 0 0	a -d c -d -a 0 0
	b -c -d 0 0 0 0	-b -c -d 0 0 0 0
(2,4,4,8)	a -b -a -a 0 -a 0	a -b -a a 0 a 0
	c -c d d 0 0 0	d -d -c -c 0 0 0
(2,4,6,12)	a -b c -a b c 0	a -b c a -b -c 0
	b a b -b d b 0	b a b b -d -b 0
(2,5,5,8)	a -d -a -a 0 -a 0	a -d -a a 0 a 0
	c -c -c -b 0 -b 0	b -b -b c 0 c 0
(2,8,8,8)	a -a b b c -d -c	a -a b b -c d c
	a a b -b c 0 c	a a b -b -c 0 -c
(3,3,3,3)	a -b -c 0 0 0 0	a b 0 -d 0 0 0
	a 0 c d 0 0 0	b -c d 0 0 0 0
(3,3,3,12)	-c -c c 0 c -a b	c c -c -d -c -a 0
	c c -c d -c 0 b	0 0 0 d 0 -a -b
(3,3,6,6)	a -d b d a 0 0	a -d c -d -a 0 0
	c d a 0 -b 0 0	c d -a 0 b 0 0
(3,4,6,8)	-c -c c 0 c -a b	c c -c 0 -c -a b
	d -b 0 b d 0 0	-d -b -a -b d 0 0
(4,4,4,4)	a -b c -d 0 0 0	a b c d 0 0 0
	a -b -c d 0 0 0	a b -c -d 0 0 0
(4,4,4,16)	c -d -c -c a -c b	c -d -c -c -a -c -b
	c -d -c c a c -b	c -d -c c -a c b
(4,4,5,5)	a -a b b 0 0 0	b -b -a -a 0 0 0
	c 0 c -d d d 0	d 0 d c -c -c 0
(4,4,8,8)	a -a b b c -d 0	a -a b b -c d 0
	a a b -b c d 0	a a b -b -c -d 0
(4,4,10,10)	b -c a -c d -d -d	b c a c -d d d
	b -d -a -d -c c c	b d -a d c -c -c

(5,5,5,5)	a -a -a -b 0 -b 0	-b b b -a 0 -a 0
	-c c c d 0 d 0	-d d d -c 0 -c 0
(6,6,6,6)	a -a b b c -d 0	-b b a a d c 0
	-c c -d -d a -b 0	-d d c c -b -a 0
(7,7,7,7)	a -a -a -b c -b d	-b b b -a d -a -c
	-c c c d a d b	-d d d -c -b -c a
(1,2,22)	a -b 0 -b b -b -b	a b 0 b -b b b
	0 b b b b -b b	c -b b b b b -b
(1,3,24)	a -c -c -c -c -c -c	b -c -c c -c c -c
	b c -c -c c -c c	b c c -c -c -c c
(1,4,20)	a -b b b b b -b	c c b -b b -b 0
	0 -b b b -b -b b	c 0 -b -b -b -b c
(1,6,12)	b b -b 0 -b -a a	-b -b b 0 b -a 0
	-b -b b 0 b 0 a	c 0 0 -a -a 0 0
(1,6,18)	c -a a -b -b a -a	-b b a a -a -a 0
	a -a a a a -b 0	a -a -a -a -b -a 0
(1,6,21)	c -b -b -b -b -b -b	a -b b -b b b -b
	a a b -b -a -b b	a -a b b -b -b b
(1,9,13)	a -b b b b b -b	0 -b b 0 b 0 0
	0 -c -c -c c c c	c -c c c -c -c c
(1,10,14)	c b -b -b -b -b b	0 b a -a a a a
	0 -b -a -b a -b a	0 -b b -a -b a b
(2,2,13)	a -b 0 0 0 0 0	a b 0 0 0 0 0
	0 -c -c -c c c c	c -c c c -c -c c
(2,7,19)	c -a a -a -a -b -a	-c -a b -a -a -a -a
	b b a -a -b -a a	b -b a a -a -a a
(2,8,13)	-b -b b -c b 0 0	-b -b b c b 0 0
	0 -a -a -a a a a	a -a a a -a -a a
(4,4,18)	a b -c b -b -b -b	a -b -c -b b b b
	a b -b -b b -c 0	a -b b b -b -c 0
(5,5,13)	a -a -a -b 0 -b 0	b -b -b a 0 a 0
	0 -c -c -c c c c	c -c c c -c -c c
(3,23)	-a 0 b -b b b b	a b b -b b b b
	a -b b b b b 0	b -b b -b b b -b
(4,19)	b -b 0 -a 0 a 0	b b a a a a a
	0 -a a 0 -a -a a	a -a a -a a a -a
(5,21)	b a a -b -b -a a	b -b a a -a -a -a
	a -a -a a -a -a 0	a -a a -a -a -a 0
(5,23)	a -a -a b b -b b	-a -b -a -b -b -b
	b -b b -b b b -b	b -b b b b -b -b
(6,17)	b b a -a a -a -a	b -b a -a -a a 0
	b -a -a -a a a a	b 0 -a 0 0 0 -a
(6,20)	b -a b -a -a 0 a	b -a -b -a 0 a -a
	b -a a a -a -a -a	b a -a a -a -a -a
(7,10)	a 0 0 -b 0 0 0	b a a 0 0 -a 0
	a 0 -b -b b 0 b	-b -b b 0 b a -a
(7,15)	b -a b -a -b 0 a	b a 0 -b 0 0 0
	b a a a 0 -a a	b a -a a a -a -a
(8,17)	a a a a -a -a 0	a -a b -b b b 0
	b a -a a b 0 a	b -a a a a -a -b
(9,16)	b -b b -a b a -b	b -a a -a -b -a 0
	b b 0 -a a a -a	a a a a -a -a 0
(9,17)	a a a a -a -a 0	a -a -a b -a a b
	a -a -b -b a -a 0	a -b -a -b b -b -b
(11,12)	a a a -a 0 0 0	-b -a a -a -a a a
	a -b a -b b b -b	b -b -b -b 0 -b 0
(11,15)	b -b -b -b 0 -b -a	b -b -b b 0 b -a
	b -a a a -a -a a	a -a a -a -a -a -a
(11,17)	b b b -b b -a -a	b b -b b b a -a
	b -a -a a -a a a	a -a a a a -a -a

(12,14) b a b -b -b -a -a b -a b -a b b a
b -b -a a a 0 -a a -a b a -b a 0