

School of Mathematics & Applied Statistics
MATH11: Mathematics Applied Mathematical Modelling 1

Assignment Week 10

__ Solutions

Spring 2004

1. The population density of fish is modelled by the differential equation

$$\frac{du}{dt} = f(u), \quad u(t=0) = u_0,$$

where the function $f(u)$ has the following properties:

- $f(0) = f(K_0) = f(K) = 0$ where $0 < K_0 < K$.
- If $u \in (0, K_0)$ then $f(u) < 0$.
- If $u \in (K_0, K)$ then $f(u) > 0$.
- If $u > K$ then $f(u) < 0$.

- (a) Sketch the growth curve $f(u)$ as a function of u .

Answer See figure 1




Figure 1: Sketch of the function described in question 1 (a).

- (b) Using your sketch determine the stability of the steady state solutions $u = 0$, $u = K_0$ and $u = K$ carefully explaining your reasoning.

Answer

The steady-state $u = 0$ is *stable*. If we change the initial condition by a small amount from $u_0 = 0$ to $u_0 = 0 + \epsilon$, where $\epsilon > 0$, the rate of change of population density is negative. Thus the solution $u(t)$ will decrease towards the steady-state solution $u = 0$.

The steady-state $u = K_0$ is *unstable*. If we change the initial condition by a small amount from $u_0 = K_0$ to $u_0 = K_0 \pm \epsilon$, where $\epsilon > 0$, the rate of change of population density is either negative ($u_0 = K_0 + \epsilon$), or positive ($u_0 = K_0 - \epsilon$). Thus the solution $u(t)$ will either decrease towards the steady-state solution $u = 0$ ($u_0 = K_0 + \epsilon$) or increase towards the solution $u = K$ ($u_0 = K_0 - \epsilon$).

The steady-state $u = K$ is *stable*. If we change the initial condition by a small amount from $u_0 = K$ to $u_0 = K \pm \epsilon$, where $\epsilon > 0$, the rate of change of population density is either negative ($u_0 = K + \epsilon$) or positive ($u_0 = K - \epsilon$). Thus the solution $u(t)$ will either decrease towards the steady-state solution $u = K$ ($u_0 = K + \epsilon$) or increase towards the steady-state solution $u = K$ ($u_0 = K - \epsilon$).

- (c) How does the long-term evolution of the differential equation depend upon the choice of the initial condition u_0 ?

Answer

- If $u_0 = 0$ then

$$u(t) = 0 \forall t.$$

- If $u_0 \in (0, K_0)$ then

$$\lim_{t \rightarrow \infty} u(t) = 0.$$

Moreover the population density decreases monotonically towards the steady-state solution $u = 0$.

- If $u_0 = K_0$ then

$$u(t) = K_0 \forall t.$$

- If $u_0 \in (K_0, K)$ then

$$\lim_{t \rightarrow \infty} u(t) = K.$$

Moreover the population density increases monotonically towards the steady-state solution $u = K$.

- If $u_0 = K$ then

$$u(t) = K \forall t.$$

- If $u_0 > K$ then

$$\lim_{t \rightarrow \infty} u(t) = K.$$

Moreover the population density decreases monotonically towards the steady-state solution $u = K$.

- (d) A disease spreads through the population reducing the population to a density $K_0/2$. What happens to the population? Justify your answer.

Answer The population will become *extinct* because the initial condition is

$$u_0 = K_0/2 \in (0, K_0).$$

The result follows from our answer to the previous question.

2. For some organisms finding a suitable mate may cause difficulties at low population density and a more realistic equation for population growth under these conditions may be

$$\frac{dN}{dt} = rN^2, \quad r > 0, N(0) = N_0.$$

- (a) Solve this problem and show that the solution becomes infinite in finite time.

Answer

$$\begin{aligned}\frac{dN}{N^2} &= r dt \\ \int_{N_0}^N \frac{dN}{N^2} &= \int_0^t r dt \\ \left[\frac{-1}{N} \right]_{N_0}^N &= rt \\ \frac{-1}{N} + \frac{1}{N_0} &= rt \\ \frac{1}{N} &= \frac{1}{N_0} - rt \\ \frac{1}{N} &= \frac{1}{N_0} - \frac{rtN_0}{N_0} \\ N &= \frac{N_0}{1 - rtN_0}.\end{aligned}$$

The solution becomes *infinite* when $t = \frac{1}{rN_0}$. \square

- (b) The model above may be improved to

$$\frac{dN}{dt} = rN^2 \left(1 - \frac{N}{K} \right).$$

Without integrating this equation find the steady-state solutions and say whether they are stable or unstable. (Do not calculate eigenvalues).

Answer The steady-state solutions of the differential equation

$$\frac{dN}{dt} = f(N)$$

are found by solving the equation

$$f(N) = 0.$$

The solutions of the equation

$$rN^2 \left(1 - \frac{N}{K} \right) = 0$$

are clearly $N = 0$ and $N = K$.

The rate of change of population density as a function of the population density is sketched in figure 2. By considering the sign of the derivative function near the two steady-state solutions we see that the steady-state solution $N = 0$ is *unstable* whilst the steady-state solution $N = K$ is *stable*. \square

- (c) Derive the model of part (b) from that of part (a). (Hint...reread Chapter 11).

Answer Consider a simple population model in which the birth and death rates are given by βN^2 and αN^2 respectively. Then the associated population model is

$$\begin{aligned}\frac{dN}{dt} &= \beta N^2 - \alpha N^2 \\ &= (\beta - \alpha) N^2 \\ &= r N^2, \quad \text{where } r = \beta - \alpha.\end{aligned}$$

This is the model analysed in part (a) of this question.




Figure 2: Sketch for question 2 (b).

In deriving this equation we have assumed that the birth rate increases with the square of the population density. However, at *high population densities* the birth rate should decrease because of competition for resources. A better assumption may be to model the birth rate by the function $\beta(1 - aN)N^2$. This leads to the differential equation

$$\begin{aligned}\frac{dN}{dt} &= \beta(1 - aN)N^2 - \alpha N^2 \\ &= [\beta(1 - aN) - \alpha]N^2 \\ &= [(\beta - \alpha) - a\beta N]N^2 \\ &= (\beta - \alpha) \left[1 - \frac{a\beta}{\beta - \alpha} N \right] N^2 \\ &= r \left(1 - \frac{N}{K} \right) N^2,\end{aligned}$$

where $r = \beta - \alpha$ and $K = \frac{\beta - \alpha}{a\beta}$.

3. Suppose a population satisfies a logistic model with carrying capacity 100 and that the population size is 10 when
- $t = 0$
- and 20 when
- $t = 1$
- . Find the intrinsic growth rate.

Use the solution to the logistic equation

$$x(t) = \frac{Kx_0}{x_0 + (K - x_0)e^{-rt}}.$$

Answer We have

$$K = 100, x_0 = 10, x(1) = 20.$$

Thus the value of r is obtained from

$$\begin{aligned} 20 &= \frac{100 \cdot 10}{10 + (100 - 10)e^{-r \cdot 1}} \\ 20 &= \frac{100 \cdot 10}{10 + 90e^{-r \cdot 1}} \\ 10 + 90e^{-r} &= 50 \\ 90e^{-r} &= 40 \\ e^{-r} &= \frac{4}{9} \\ \frac{9}{4} &= e^r \\ r &= \ln \frac{9}{4} \\ &\approx 0.8109 \quad (4 \text{ dp}) \end{aligned}$$

□

4. Nisbet & Gurney (1983) suggested the following form for the per-capita growth rate

$$r(x) = r \exp \left[1 - \frac{x}{K} \right] - d$$

Consider the associated population model

$$\frac{dx}{dt} = \left(r \exp \left[1 - \frac{x}{K} \right] - d \right) x, \quad x(0) = x_0, r > de^{-1}.$$

(a) Find the steady-state(s) of the model. How do the number of steady-state solutions ($x^* \geq 0$) and their biological feasibility depend upon the values of r and d ?

Answer The steady-state solutions for the equation

$$\frac{dx}{dt} = f(x)$$

are the values of x, x^* , for which

$$\begin{aligned} f(x) &= 0. \\ \text{i.e. } \left(r \exp \left[1 - \frac{x}{K} \right] - d \right) x &= 0. \end{aligned}$$

The solutions of this equation are given by

$$x^* = 0$$

and

$$\begin{aligned} r \exp \left[1 - \frac{x^*}{K} \right] - d &= 0 \\ \Rightarrow \exp \left[1 - \frac{x^*}{K} \right] &= \frac{d}{r} \\ \Rightarrow 1 - \frac{x^*}{K} &= \ln \left(\frac{d}{r} \right) \\ \Rightarrow \frac{x^*}{K} &= 1 - \ln \left(\frac{d}{r} \right) \\ \Rightarrow x^* &= K \left[1 - \ln \left(\frac{d}{r} \right) \right] \end{aligned}$$

There are *two* steady-state solutions: $x^* = 0$, and $x^* = K \left[1 - \ln \left(\frac{d}{r} \right) \right]$. The second of these is biologically feasible if

$$\begin{aligned} K \left[1 - \ln \left(\frac{d}{r} \right) \right] &> 0 \\ \Rightarrow 1 &> \ln \left(\frac{d}{r} \right) \\ \Rightarrow e &> \frac{d}{r} \end{aligned}$$

(b) Explain why it is reasonable to assume that $r > de^{-1}$.

If $de^{-1} > r$ then there is only one biologically feasible steady-state solution, $x^* = 0$. A graph of function $f(x)$ shows that in this case the population becomes extinct. This is not very interesting. So, let us suppose that $de^{-1} < r$.

(c) Sketch $\frac{dx}{dt}$ as a function of x . Hence determine how the long-term dynamics of the model depend upon the initial value x_0 .

Answer The function is sketched in figure 3.




Figure 3: Sketch of the rate of change of population density for the Nisbet & Gurney model.

Let $x_1^* = 0$ and $x_2^* = K \left[1 - \ln \left(\frac{d}{r} \right) \right]$.

- (i) If $x(0) = 0$ then $x(t) = 0 \quad \forall t$.
- If $x(0) = x_2^*$ then $x(t) = x_2^* \quad \forall t$.
- (ii) If $x(0) \in (0, x_2^*)$ then $\lim_{t \rightarrow \infty} x(t) = x_2^*$.

(iii) If $x(0) > x_2^*$ then $\lim_{t \rightarrow \infty} x(t) = x_2^*$.

□