

If  $f(z) = 2z$  and  $g(x) = \frac{1}{x^2}$  what is  $(g(f(z)))$ .

(a)  $\frac{2z}{x^2}$ .

(b)  $\frac{2}{z^2}$ .

(c)  $\frac{2}{x^2}$ .

(d)  $\frac{1}{4x^2}$ .

(e)  $\frac{1}{4z^2}$ .

If  $f(z) = 2z + 1$  and  $g(x) = \frac{1}{x}$  what is  $g(f(a))$ ?

1.  $\frac{2}{x} + 1$

2.  $\frac{2}{a} + 1$

3.  $\frac{1}{2x + 1}$

4.  $\frac{1}{1 + 2a}$

5.  $\frac{1}{2a} + 1$

**answer**

If  $f(z) = 2z + 1$  and  $g(x) = \frac{1}{x}$  then  $f(g(a))$  is given by

(a)  $\frac{2}{a} + 1.$

(b)  $\frac{2}{x} + 1.$

(c)  $\frac{1}{2a + 1}.$

(d)  $\frac{1}{2x + 1}.$

(e)  $\frac{1}{2a} + 1.$

**answer**

$\log \left( x + \sqrt{1 + x^2} \right)$  equals

(a)  $\log x + \frac{1}{2} \log (1 + x^2)$

(b)  $\frac{1}{2} \log x \cdot \log (1 + x^2)$

(c) neither

$\frac{d}{dx} \cosh (x^2 + 1)$  equals

(a)  $\sinh (x^2 + 1)$

(b)  $2x \sinh (x^2 + 1)$

(c)  $-\sinh (x^2 + 1)$

(d)  $-2x \sinh (x^2 + 1)$

(e)  $\sinh x^3 + x$

Match the items in column 1 with the definitions in column 2.

- |                                |   |
|--------------------------------|---|
| (a) $\sinh x$                  | (A) $(e^x - e^{-x}) / 2$                    |
| (b) $\cosh x$                  | (B) 1                                       |
| (c) $\cosh^2 x + \sinh^2 x$    | (C) $1 + 2 \sinh^2 x$                       |
| (d) $\operatorname{arccosh} x$ | (D) $(e^x + e^{-x}) / 2$                    |
| (e) $\cosh^2 x - \sinh^2 x$    | (E) $\ln \left( x + \sqrt{x^2 - 1} \right)$ |



$\frac{d}{dx}(-2xy)$  equals

(a)  $-2y'$

(b)  $-2xy' + 2y$

(c)  $-2xy' - 2y$

$\frac{d}{dx} (yy')$  equals

(a)  $y'y''$

(b)  $y' + yy''$

(c)  $(y')^2 + yy''$