

Fundamentals

Lecture Two

Indices Simplify

$$6x^3y^{-2} \times \frac{1}{24}x^{-5}y^4$$

Surds Simplify

1. $\frac{\sqrt{5}}{\sqrt{45}}$

2. $\frac{5}{1+\sqrt{5}}$

Definition

If a is a real number and n is a natural number, then

$$a^n = a \times a \times \dots \times a \quad (n \text{ factors}).$$

n is called either the *index* of a , the *exponent* on a , or the *power* to which a is raised. a is called the *base*. \square

Rules

$$1. \quad a^m \times a^n = a^{m+n}$$

$$2. \quad a^m \div a^n = a^{m-n}$$

$$3. \quad (ab)^m = a^m b^m$$

$$4. \quad (a^m)^n = a^{mn}$$

$$5. \quad a^{-m} = \frac{1}{a^m}$$

$$6. \quad a^0 = 1$$

$$7. \quad a^{m/n} = ({}^n\sqrt{a})^m = {}^n\sqrt{a^m}$$

Examples

Simplify the following expressions

1.
$$\frac{3xy^2 \times 4x^3 \times y}{2xy \times 4y^2}$$

2.
$$\frac{6x^{-4} \times 2x^3}{3x^{-3}}$$

3.
$$(5x^2y^{-3/2}z^{1/4})^2 \times (4x^4y^2z)^{-1/2}$$

$$\left(\frac{3}{2}x^3\right)$$

$$(4x^2)$$

$$\left(\frac{25}{2}x^2y^{-4}\right)$$

Definition

A *surd* is the name given to an *irrational* number that can be expressed in the form ${}^n\sqrt{a}$.

(See definition of irrational p1-1)

Notation

\sqrt{a} , \sqrt{a} or $a^{1/2}$ means “square root of a ”

${}^n\sqrt{a}$, ${}^n\sqrt{a}$ or $a^{1/n}$ means “ n^{th} root of a ”

Irrational numbers such as $\sqrt{2}$, $5\sqrt{3}$, ${}^3\sqrt{7}$... are *surds*, whereas $\sqrt{4}$, ${}^3\sqrt{27}$, $3\sqrt{9}$... are *not*, since they can be evaluated *exactly*.

Question. Evaluate exactly $\sqrt{4}$, ${}^3\sqrt{27}$ & $3\sqrt{9}$.

Rules

$$1. \sqrt{xy} = \sqrt{x} \times \sqrt{y}$$

$$2. \sqrt{\frac{x}{y}} = \frac{\sqrt{x}}{\sqrt{y}}$$

$$3. (\sqrt{x})^2 = x \text{ provided } x \geq 0$$

$$4. \sqrt{x^2} = \begin{cases} +x & \text{if } x \geq 0, \\ -x & \text{if } x < 0. \end{cases}$$

$$5. a\sqrt{x} \pm b\sqrt{x} = (a \pm b)\sqrt{x}$$

Examples

Simplify the following expressions

1. $4\sqrt{3} + 5\sqrt{7} - 2\sqrt{3} + \sqrt{7}$

2. $2\sqrt{3} \times 4\sqrt{5}$

3. $\sqrt{18} \div \sqrt{2}$

4. $6\sqrt{15} \div 2\sqrt{3}$

5. $\sqrt{8}$

6. $\sqrt{3} \times \sqrt{21}$

7. $5\sqrt{18}$

8. $\sqrt{20} + \sqrt{45}$

9. $\sqrt{12} - \sqrt{27} + \sqrt{75} - \sqrt{15}$

$$(2\sqrt{3} + 6\sqrt{7})$$

$$(8\sqrt{15})$$

$$(\pm 3)$$

$$(3\sqrt{5})$$

$$(2\sqrt{2})$$

$$(3\sqrt{7})$$

$$(15\sqrt{2})$$

$$(5\sqrt{5})$$

$$(\sqrt{3} [4 - \sqrt{5}])$$

Rationalising the Denominator

The *conjugate* of $\sqrt{a} + \sqrt{b}$ is $\sqrt{a} - \sqrt{b}$.

Multiplying a surd by its conjugate always gives a rational number

$$\begin{aligned}(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b}) \\ &= (\sqrt{a})^2 - (\sqrt{b})^2, \\ &= a - b.\end{aligned}$$

Using this rule we can express any surd with a rational denominator.

Examples

Simplify the following expressions.

1. $\frac{1}{\sqrt{a}}$

2. $\frac{1}{\sqrt{a} + \sqrt{b}}$

3. $\frac{3}{\sqrt{5}}$

4. $\frac{2}{6 - \sqrt{3}}$

5. $\frac{\sqrt{3} + \sqrt{2}}{\sqrt{3} - \sqrt{2}}$

$$\left(\frac{\sqrt{a}}{a}\right)$$

$$\left(\frac{\sqrt{a}-\sqrt{b}}{a-b}\right)$$

$$\left(\frac{3\sqrt{5}}{5}\right)$$

$$\left(\frac{12+2\sqrt{3}}{33}\right)$$

$$(5 + 2\sqrt{6})$$

Equality of surds

If $a + \sqrt{b} = c + \sqrt{d}$ where a and c are rational and \sqrt{b} and \sqrt{d} are surds, then $a = c$ and $b = d$.

Examples

Find the values of α and β in the following expressions

$$1. \alpha + \sqrt{\beta} = 2 + \sqrt{5}$$

$$2. 2\alpha + 3\sqrt{\beta} = 4 + 3\sqrt{2}$$

$$3. \alpha\sqrt{3} = \sqrt{27}$$

$$4. \alpha + 2\sqrt{\beta} = (\sqrt{3} + \sqrt{5})^2$$

$$(\alpha = 2, \beta = 5)$$

$$(\alpha = 2, \beta = 2)$$

$$(\alpha = 3)$$

$$(\alpha = 8, \beta = 15)$$

Exercises on Indices and Surds

Exercise 1.2.4 (pages 4& 5)

For each of the questions do as many of the subquestions as you require in order to gain mastery of the basic technique.

DO NOT do question 4 on the surds worksheet.